

Comprehensive evaluation of abiotic stress tolerance and graft compatibility of *Citrus junos* cv. 'Shuzhen No.1'

Wen He^{1,2#}, Rui Xie^{1#}, Jiufeng Chai¹, Hao Wang¹, Yan Wang^{1,2}, Qing Chen^{1,2}, Zhiwei Wu¹, Mengyao Li¹, Yuanxiu Lin¹, Yunting Zhang¹, Ya Luo¹, Yong Zhang¹, Haoru Tang¹ and Xiaorong Wang^{1,2*}

¹ College of Horticulture, Sichuan Agricultural University, Chengdu 611130, Sichuan, PR China

² Key Laboratory of Agricultural Bioinformatics, Ministry of Education, Chengdu 611130, PR China

These authors contributed equally: Wen He, Rui Xie

* Corresponding author, E-mail: wangxr@sicau.edu.cn

Abstract

Citrus is one of the world's most economically important fruit crops cultivated by grafting. To support the growth of scion cultivars, rootstock is the primary source of resistance to various abiotic stresses. Herein, seedlings of two genotypes of *Citrus junos* Sieb. ex Tanaka (the novel rootstock 'Shuzhen No.1' and commonly used rootstock 'Ziyang Xiangcheng'), as well as three commonly used rootstocks including citrange (*Citrus sinensis* Osbeck. × *Poncirus trifoliata* Raf.), trifoliolate orange (*P. trifoliata*), and red tangerine (*Citrus tangerine* Hort. Ex Tanaka), were used as testing materials. The seed characteristics were evaluated, and the rootstock seedlings were subjected to flooding, drought, alkaline, and freezing treatments. Over time, the contents of chlorophyll, soluble sugar, proline, malondialdehyde, and the activity of superoxide dismutase, peroxidase, and catalase in the leaves under different treatments were examined. Furthermore, five citrus varieties were grafted as scions onto one-year-old seedlings from the four rootstocks. Graft success, shoot growth, and leaf greenness were measured and compared. The physiological and biochemical changes in 'Shuzhen No.1' were found to be similar to those in 'Ziyang Xiangcheng'. 'Shuzhen No.1' exhibited greater tolerance to flooding, alkaline, and freezing stress compared to the other four widely used citrus rootstocks, as indicated by physiological and biochemical indexes and principal component analysis. Moreover, the five citrus varieties grafted onto 'Shuzhen No.1' demonstrated vigorous growth and tree vigor. These findings provide valuable insights for the application of 'Shuzhen No.1' and future research on citrus rootstock.

Citation: He W, Xie R, Chai J, Wang H, Wang Y, et al. 2024. Comprehensive evaluation of abiotic stress tolerance and graft compatibility of *Citrus junos* cv. 'Shuzhen No.1'. *Fruit Research* 4: e006 <https://doi.org/10.48130/frures-0023-0042>

Introduction

Citrus is the world's most economically important fruit crop, and the majority of citrus is grown in mountainous regions with barren soil^[1]. Citrus productivity can be greatly affected by environmental changes, such as brief periods of flooding, drought, or cold^[2–4]. Moreover, citrus is susceptible to alkaline soils and exhibits leaf/shoot chlorosis, limiting its geographical distribution^[5,6]. Citrus is cultivated through grafting, and rootstock can modify scion architecture and act as the core source of resistance to various stresses, allowing the upper section growth of scion cultivars to thrive^[7,8]. With growing interest in perennial crops as valuable components of sustainable agriculture, rootstocks provide an approach for improving and expanding citrus perennial cultivation under various environmental conditions^[9].

Generally, rootstock selection and use are mainly determined by compatibility, orchard soil conditions, and local citriculture practice^[10]. Germplasm with polyembryonicity, which can develop one or more somatic embryos that are genetically identical to the mother tree, is often selected for citrus rootstock due to genetically uniform rootstocks which can feasibly be prepared solely by sowing seeds^[11,12]. Many rootstocks are used in citrus cultivation. Trifoliolate orange (*Poncirus trifoliata* (L.) Raf) is widely used in citrus breeding owing to its cold

hardiness and disease resistance^[13]. However, trifoliolate orange is sensitive to alkalinity and mineral deficiency and is incompatible with some citrus cultivars^[5,14–16]. Citrange (*Citrus sinensis* × *P. trifoliata*) is drought tolerant yet susceptible to salt and alkalinity^[17,18]. Red tangerine (*Citrus reticulata* Blanco) is resistant to B-deficiency and citrus exocortis viroid (CEVd); however, the fruit quality of scion degrades when red tangerine is used as a rootstock^[19]. *Citrus junos* Sieb. Ex Tanaka is an iron-deficient, alkaline-, cold- and acid-tolerant citrus rootstock native to southwest China^[5,17,20]. Abiotic stresses can alter osmotic equilibrium and induce oxidative stress in plants through excessive generation of reactive oxygen species (ROS)^[21,22]. Plants neutralize these ROS through different mechanisms, which can be classified as non-enzymatic and enzymatic antioxidant systems, including antioxidant enzymes superoxide dismutase (SOD), peroxidases (POD), and catalase (CAT)^[23–25]. Tolerant species or genotypes exhibit higher antioxidant enzyme activities than sensitive genotypes. Investigating tolerance to different abiotic stresses is critical for identifying the genetic resource for abiotic stress tolerance. Although rootstock can influence the agronomic performance of citrus trees, some widely used rootstocks may still demonstrate graft incompatibility in the orchard^[9,10]. Graft compatibility of intergeneric and intrafamilial species represents a tremendous agronomic

potential for genetic improvement and improved crop management by combining unique traits from wild relative rootstocks with commercial citrus scion varieties^[7].

In our previous citrus rootstock breeding effort, we reported a novel rootstock cultivar, *C. junos* cv. Shuzhen No.1, with vigorous growth, spherical crown, upright and dense hard branches, cold resistance, and robust adaptation to basic soil conditions^[26,27]. Therefore, this study compared the differences in seed germination characteristics, abiotic tolerance (drought, flooding, alkaline, and freezing), and grafted plant performance between 'Shuzhen No.1' and other common citrus rootstocks. This study determined the polyembryony and seedling emergence traits of five citrus rootstocks, comprehensively evaluated the tolerance responses of different genotypes of citrus rootstock, and provided information on the performances of five citrus varieties on rootstocks. Our findings provide insights into rootstock selection and promote the utilization of the new citrus rootstock 'Shuzhen No.1'.

Materials and methods

Plant materials and culture conditions

Mature fruits of 'Shuzhen No.1' (abbreviated CjSz), trifoliolate orange (Pt), red tangerine (Ct), citrange (Cp), and 'Ziyang Xiangcheng' (CjZy) were harvested for collecting seeds from the Citrus Germplasm Repository of Sichuan Province, Chengdu, China. Isolated seeds were surface sterilized using 0.5 M NaOH as described previously^[17]. Uniform-sized seedlings were selected and grown in a growth chamber in perlite-filled pots. All seedlings were cultured for approximately six months with normal watering and fertilization.

Determination of polyembryony and seedling emergence traits

Two hundred viable seeds (not replicated) were selected from each rootstock, and the following parameters were assessed: (1) cumulative seedling number, (2) percentage of single seed emergence, and (3) polyembryony. *CitRWP* plays a principal role in regulating somatic embryogenesis in citrus nucellar tissues, and its alleles were divided into two types and polyembryonic alleles with a MITE insertion^[11,12]. The absence or presence of the MITE insertion was evaluated with genomic PCR using the following primer set: forward 5'-GTTACTTGGA GACGGCCTAACG-3' and reverse 5'-TCGATCATGTAATGCTGACT C-3'^[11].

Stress treatments

Abiotic stress treatments included flooding (roots submerged in water with only stem and leaves exposed to the environment for 6 and 7 weeks), drought (20% soil water content for 1 and 2 weeks), alkalinity (watering distilled water with pH 8.0 and 9.0 for 8 weeks), and freezing (−10 °C for 1 and 2 h) were performed. Three biological replicates (five seedlings per replicate) were set randomly for each treatment. All leaves were sampled from five rootstocks, frozen in liquid nitrogen, and stored at −80 °C.

Determination of physiological and biochemical indexes

The total chlorophyll and carotenoid contents were measured using the method reported by Lichtenthaler & Buschmann^[28]. Fresh leaves (1 g) were ground in a freezing

mortar and pestle with 10 mL of 80% acetone. Following filtering, the pigment solution's optical density (OD) was measured at 470, 645, and 663 nm to determine carotenoid, chlorophyll (Chl) *a*, Chl *b*, and total Chl content, respectively. The assessed photosynthetic pigments were presented in mg/g fresh weight (FW). Antioxidant enzyme activities of SOD (EC 1.15.1.1), guaiacol peroxidase (POD, EC 1.11.1.7), and CAT (EC 1.11.1.6) were determined as previously described^[29]. Malondialdehyde (MDA) content was measured using the thiobarbituric acid (TBA) method^[30]. Soluble sugars and proteins were analyzed as previously described^[31]. To minimize the differences between different genotypes, the data were expressed as ratios relative to the values of control groups.

Performances of different graft combinations

'Chunjian' (*C. reticulata* × (*C. reticulata* × *C. sinensis*)), 'Buzhi-huo' (*C. unshiu* × *C. sinensis*), 'Mingrijian' ((*C. unshiu* × *C. hassaku*) × *C. sinensis*), 'Dafen' (*C. unshiu*), and 'Tarocco' (*C. sinensis*) were grafted onto four rootstocks, including CjSz, CjZy, Ct, and Pt. Ninety seedlings were cultured for 1 month with normal watering and fertilization, and their survival rates were measured on March 30th, 2019. Tree growth and leaf greenness were assessed in 5–10 grafted trees in each cultivar. Stem thicknesses below and above the graft joint were measured using a vernier caliper on October 30th, 2020. Shoot length and longitudinal and horizontal growth of trees were recorded using a tape measure from summer shoots on October 30th, 2020. Ten mature leaves from summer shoots were selected from each tree to measure soil-plant analysis development (SPAD) with a SPAD-502 chlorophyll meter.

Statistical analysis

Microsoft Excel was used to prepare the collected data. Significant differences between grafted combinations were analyzed using Tukey's method, and Pearson correlation and principal component analyses were performed using SPSS 20.0 software. All figures were drawn using GraphPad Prism (v. 7.04).

Results

Polyembryony and seedling emergence traits among tested rootstocks

Compared with CjZy, CjSz has a spherical crown and upright and dense hard branches (Fig. 1a, b). CjSz and CjZy had solitary flowers (Fig. 1c, d). However, the lateral petals of CjSz are purple from bud to bloom, with a flower diameter of 2.0 cm (Fig. 1c). The fruits of both CjSz and CjZy were orange in color at maturity (Fig. 1c, d) and both were polyembryonic (Fig. 1c, d). Citrus rootstock seeds germinated 40–45 d after seeding and stopped germinating 95–100 d later (Fig. 1e). CjSz had the greatest emergence rate and polyembryonic ratio among the tested rootstocks, reaching 160.77% and 70.69%, respectively (Fig. 1e, f). The germination potential was 25.50%, which was lower than that of trifoliolate orange (29.00%) but higher than that of CjZy, citrange, and red tangerine (Fig. 1e). According to statistics on single seed emergences, CjSz had the most single seed emergences (two, up to 44.83%), slightly lower than CjZy (46.76%) but higher than Pt, Ct, and Cp. The proportion of CjSz was highest among the five tested rootstocks, at 18.72% (three seedlings), 4.39% (four seedlings), 1.48% (five seedlings), and 0.49% (six seedlings). Additionally, the maximum number of seedlings per grain was seven (Fig. 1f). These results are




Fig. 1 Comparison of morphology between two genotypes of *Citrus junos*. (a), (b) Six year old trees; (c), (d) flowers, fruits and seeds; (e) cumulative number of seedlings; (f) percentage of single seed emergence; (g) MITE insertion in five rootstock germplasms. CjSz: Shuzhen No.1 (*Citrus junos* Sieb. Tanaka); CjZy: Ziyang Xiangcheng (*C. junos* Sieb. Tanaka); Cp: citrange (*C. sinensis* Osbeck. × *Poncirus trifoliata* Raf.); Pt: trifoliolate orange (*P. trifoliata* [L.] Raf) and Ct: Red tangerine (*C. tangerine* Hort. Ex Tanaka). Scale bars = 1 cm.

consistent with the MITE insertion detection results (Fig. 1g). Apomixis in citrus is sporophytic and highly stable across commercial varieties. *Citrus junos* fruits were densely seeded, with most of the seeds being plump and polyembryonic, which can generate large numbers of uniform rootstocks from seeds^[12,31].

Leaf photosynthetic pigments under abiotic stresses

All genotypes developed leaf chlorosis at the end of the abiotic stress treatments (Fig. 2). Almost all treatments reduced the content of leaf photosynthetic pigments in all rootstocks (Table 1). Comparing the pigment content data among the different rootstocks, it is evident that CjSz demonstrated better adaptability to alkaline and freezing stresses (Table 1, Fig. 2). Specifically, under alkaline treatment with a pH of 8.0, CjSz exhibited the smallest decrease in leaf photosynthetic pigments content. In the case of alkaline treatment with a pH of 9.0, CjSz displayed similar levels of leaf chlorina compared to Cp, followed by CjZy, Pt and Ct. Similarly, under freezing treatment for 1 h, the ratios of Chl a, total Chl and total carotenoids in CjSz were higher than in other rootstocks, although the differences were not statistically significant. More specifically, among the different treatments, CjSz experienced the greatest decrease in Chl a, Chl b, and total Chl under 2 weeks of drought treatment, with ratios of 0.48, 0.64, and 0.53, respectively.

Conversely, CjSz demonstrated the least decrease under freezing treatment for 1 h, with ratios of 0.96, 1.05, and 0.99, respectively (Table 1). The total carotenoid content in CjSz experienced the most significant decrease after 2 weeks of drought treatment, while the least decrease occurred under alkaline treatment with a pH of 8.0 (Table 1).

MDA, soluble protein, and soluble sugar contents and SOD, POD, and CAT activities under abiotic stresses

The levels of MDA, and the activities of SOD, POD, and CAT were significantly influenced by abiotic stresses (Table 2). CAT activity decreased under flooding, alkaline, and freezing stress, but slightly increased under drought stress. MDA levels and SOD and POD activities increased under flooding, drought, alkaline, and freezing stresses in all citrus rootstock genotypes (Table 2). However, CAT activity decreased under abiotic stress. Among the genotypes, CjSz exhibited the highest increase in MDA levels during 7 weeks of flooding treatment, and the lowest increase during 1 h of freezing treatment. SOD activity in CjSz showed the greatest increase after 7 weeks of flooding treatment and the smallest increase after alkaline stress treatment at pH 8.0. CjSz had the highest POD activity ratio of 1.62 under freezing stress, and the lowest value of 1.25 under 2 weeks of drought treatment. In comparison to the other four rootstocks, CjSz had the highest SOD ratio value (1.71) under 7




Fig. 2 Growth state of five rootstocks under abiotic stresses. (a) Flooding stress; (b) drought stress; (c) alkaline stress (pH = 8.0 and pH = 9.0); (d) freezing stress.

Table 1. Statistics for chlorophyll a, chlorophyll b, total chlorophyll and total carotenoid compared with controls.

| Species | Flooding stress | | Drought stress | | Alkaline stress | | Freezing stress | | |
|---------|-----------------|---------------|----------------|--------------|-----------------|---------------|-----------------|---------------|---------------|
| | 6 weeks | 7 weeks | 1 week | 2 weeks | pH = 8.0 | pH = 9.0 | 1 h | 2 h | |
| Chl a | CjSz | 0.72 ± 0.12a | 0.60 ± 0.04a | 0.53 ± 0.08a | 0.48 ± 0.15a | 0.86 ± 0.26a | 0.74 ± 0.35ab | 0.96 ± 0.24a | 0.83 ± 0.11a |
| | CjZy | 0.67 ± 0.22a | 0.68 ± 0.09a | 0.59 ± 0.16a | 0.45 ± 0.22a | 0.79 ± 0.14a | 0.69 ± 0.19ab | 0.85 ± 0.17a | 0.79 ± 0.19a |
| | Cp | 0.77 ± 0.26a | 0.69 ± 0.26a | 0.66 ± 0.28a | 0.61 ± 0.16a | 0.83 ± 0.26a | 0.77 ± 0.20a | 0.88 ± 0.12a | 0.85 ± 0.18a |
| | Pt | 0.63 ± 0.06ab | 0.63 ± 0.19a | 0.63 ± 0.03a | 0.60 ± 0.08a | 0.72 ± 0.03ab | 0.68 ± 0.07ab | 0.91 ± 0.20a | 0.88 ± 0.05a |
| | Ct | 0.44 ± 0.10b | 0.38 ± 0.09b | 0.28 ± 0.05b | 0.23 ± 0.07b | 0.48 ± 0.04b | 0.47 ± 0.07b | 0.92 ± 0.06a | 0.69 ± 0.15a |
| Chl b | CjSz | 0.78 ± 0.12a | 0.70 ± 0.08a | 0.67 ± 0.14a | 0.64 ± 0.15a | 0.87 ± 0.26a | 0.77 ± 0.31a | 1.05 ± 0.22a | 1.00 ± 0.12a |
| | CjZy | 0.70 ± 0.20a | 0.65 ± 0.09a | 0.72 ± 0.09a | 0.52 ± 0.20a | 0.76 ± 0.11a | 0.74 ± 0.13a | 0.86 ± 0.11ab | 0.92 ± 0.17ab |
| | Cp | 0.76 ± 0.24a | 0.68 ± 0.25a | 0.71 ± 0.25a | 0.68 ± 0.15a | 0.85 ± 0.18a | 0.78 ± 0.16a | 0.82 ± 0.09b | 0.83 ± 0.11b |
| | Pt | 0.71 ± 0.13a | 0.66 ± 0.20a | 0.66 ± 0.05a | 0.68 ± 0.11a | 0.69 ± 0.09ab | 0.66 ± 0.08ab | 0.86 ± 0.19ab | 0.87 ± 0.11ab |
| | Ct | 0.47 ± 0.09b | 0.50 ± 0.12a | 0.39 ± 0.04b | 0.33 ± 0.06b | 0.51 ± 0.04b | 0.47 ± 0.05b | 1.00 ± 0.10ab | 0.64 ± 0.12c |
| Total | CjSz | 0.74 ± 0.12a | 0.63 ± 0.05ab | 0.57 ± 0.10a | 0.53 ± 0.15a | 0.86 ± 0.26a | 0.75 ± 0.33a | 0.99 ± 0.23a | 0.89 ± 0.12a |
| Chl | CjZy | 0.68 ± 0.21a | 0.67 ± 0.09a | 0.63 ± 0.14a | 0.47 ± 0.21a | 0.78 ± 0.13a | 0.71 ± 0.17ab | 0.85 ± 0.15a | 0.84 ± 0.19ab |
| | Cp | 0.76 ± 0.25a | 0.68 ± 0.26a | 0.67 ± 0.27a | 0.63 ± 0.16a | 0.84 ± 0.23a | 0.77 ± 0.19a | 0.86 ± 0.11a | 0.84 ± 0.15ab |
| | Pt | 0.65 ± 0.08ab | 0.64 ± 0.19a | 0.63 ± 0.03a | 0.63 ± 0.09a | 0.71 ± 0.05ab | 0.67 ± 0.07ab | 0.89 ± 0.19a | 0.87 ± 0.07a |
| | Ct | 0.45 ± 0.10b | 0.42 ± 0.10b | 0.32 ± 0.04b | 0.27 ± 0.07b | 0.49 ± 0.03b | 0.47 ± 0.06b | 0.95 ± 0.08a | 0.67 ± 0.14b |
| | Total | CjSz | 0.83 ± 0.12a | 0.65 ± 0.10a | 0.68 ± 0.11a | 0.55 ± 0.18ab | 0.96 ± 0.30a | 0.71 ± 0.27a | 0.96 ± 0.20a |
| Car | CjZy | 0.62 ± 0.15bc | 0.65 ± 0.08a | 0.56 ± 0.14a | 0.51 ± 0.25ab | 0.68 ± 0.11bc | 0.69 ± 0.12ab | 0.77 ± 0.14a | 0.76 ± 0.12bc |
| | Cp | 0.76 ± 0.24ab | 0.67 ± 0.23a | 0.63 ± 0.19a | 0.69 ± 0.13a | 0.79 ± 0.15ab | 0.75 ± 0.17a | 0.92 ± 0.06a | 0.86 ± 0.14ab |
| | Pt | 0.69 ± 0.08ab | 0.62 ± 0.16ab | 0.64 ± 0.04a | 0.65 ± 0.09a | 0.73 ± 0.04b | 0.63 ± 0.05ab | 0.91 ± 0.23a | 0.95 ± 0.05a |
| | Ct | 0.45 ± 0.12c | 0.43 ± 0.13b | 0.38 ± 0.05b | 0.37 ± 0.10b | 0.47 ± 0.03c | 0.49 ± 0.05b | 0.89 ± 0.14a | 0.67 ± 0.14c |

Note: Chl: chlorophyll, Car: carotenoids. Data shown in the table are expressed as ratios relative to the values obtained on control seedlings. Three biological replicates (five seedlings per replicate) were set randomly for each treatment. Significance was tested for indicators of different rootstocks in the same treatment, and different lowercase letters indicate significant differences at $p < 0.05$.

weeks of flooding stress. Additionally, CjSz exhibited the highest POD ratio value under alkaline (1.37 and 1.46) and freezing stresses (1.62). Under drought stress, CjSz had significantly lower MDA and POD ratio values than the other four rootstocks.

The regulation of soluble protein and sugar contents in response to abiotic stressors varied among the citrus rootstock genotypes (Table 3). Compared with other rootstocks, CjSz had the lowest ratio of soluble proteins during 7 weeks of flooding stress, but the highest ratio under drought, alkaline, and freezing stresses. CjSz exhibited the greatest increase in soluble sugars under 7 weeks of flooding stress and 1 week of drought

stress, whereas other rootstocks showed moderate increases. Specifically, the overall ratio of soluble proteins in CjSz was the highest (2.16) under 2 h of freezing stress and the lowest (1.10) during 7 weeks of flooding stress. However, the ratio of soluble sugars displayed opposite trends.

Comprehensive evaluation of rootstocks under abiotic stresses

To provide a comprehensive assessment of the tolerance of different rootstocks, we conducted a principal component analysis to calculate various physiological and biochemical

Table 2. Mean comparison of malondialdehyde (MDA), superoxide dismutase (SOD), catalase (CAT) and peroxidase (POD).

| Species | Flooding stress | | Drought stress | | Alkaline stress | | Freezing stress | | |
|---------|-----------------|----------------|----------------|---------------|-----------------|---------------|-----------------|----------------|---------------|
| | 6 weeks | 7 weeks | 1 week | 2 weeks | pH = 8.0 | pH = 9.0 | 1 h | 2 h | |
| MDA | CjSz | 1.42 ± 0.30bc | 2.02 ± 0.16ab | 1.07 ± 0.11d | 1.90 ± 0.26b | 1.11 ± 0.11b | 1.50 ± 0.13c | 1.02 ± 0.22b | 1.24 ± 0.39b |
| | CjZy | 1.45 ± 0.17bc | 1.81 ± 0.30b | 1.79 ± 0.22b | 2.10 ± 0.14a | 1.47 ± 0.30ab | 1.75 ± 0.28bc | 1.02 ± 0.13b | 1.15 ± 0.27b |
| | Cp | 1.67 ± 0.20ab | 1.84 ± 0.41b | 1.31 ± 0.23cd | 2.25 ± 0.61a | 1.33 ± 0.39ab | 1.36 ± 0.20c | 1.49 ± 0.29a | 1.92 ± 0.63a |
| | Pt | 1.26 ± 0.16c | 1.48 ± 0.34b | 1.45 ± 0.31bc | 2.03 ± 0.43a | 1.84 ± 0.40a | 2.41 ± 0.42a | 0.91 ± 0.34b | 1.01 ± 0.54b |
| | Ct | 1.75 ± 0.25a | 2.42 ± 0.66a | 2.30 ± 0.36a | 2.29 ± 0.37a | 1.36 ± 0.10ab | 1.92 ± 0.36b | 1.00 ± 0.17b | 1.46 ± 0.26ab |
| SOD | CjSz | 1.60 ± 0.26a | 1.71 ± 0.03a | 1.53 ± 0.02ab | 1.63 ± 0.11a | 1.25 ± 0.08a | 1.31 ± 0.15ab | 1.31 ± 0.06ab | 1.35 ± 0.16a |
| | CjZy | 1.65 ± 0.10a | 1.70 ± 0.07a | 1.62 ± 0.07a | 1.71 ± 0.06a | 1.31 ± 0.29a | 1.36 ± 0.29a | 1.12 ± 0.27b | 1.54 ± 0.21a |
| | Cp | 1.25 ± 0.18c | 1.19 ± 0.18b | 1.18 ± 0.16c | 1.15 ± 0.15b | 1.12 ± 0.05ab | 1.26 ± 0.02ab | 1.25 ± 0.17ab | 1.05 ± 0.23b |
| | Pt | 1.56 ± 0.10ab | 1.50 ± 0.46ab | 1.69 ± 0.16a | 1.51 ± 0.28a | 0.94 ± 0.30b | 0.94 ± 0.39c | 1.39 ± 0.19a | 1.56 ± 0.11a |
| | Ct | 1.36 ± 0.10bc | 1.61 ± 0.18a | 1.35 ± 0.17bc | 1.23 ± 0.18b | 1.08 ± 0.07ab | 1.04 ± 0.09bc | 1.18 ± 0.24ab | 1.07 ± 0.08b |
| POD | CjSz | 1.40 ± 0.31a | 1.50 ± 0.10ab | 1.25 ± 0.38c | 1.45 ± 0.61b | 1.37 ± 0.49a | 1.46 ± 0.22a | 1.62 ± 0.32ab | 1.62 ± 0.37a |
| | CjZy | 1.61 ± 0.33a | 1.62 ± 0.38ab | 2.23 ± 0.68ab | 2.69 ± 0.72a | 1.29 ± 0.24a | 1.38 ± 0.54a | 1.25 ± 0.66b | 1.55 ± 0.59a |
| | Cp | 1.45 ± 0.16a | 2.16 ± 0.86a | 2.59 ± 1.09a | 2.90 ± 0.70a | 1.27 ± 0.28a | 1.31 ± 0.32a | 1.28 ± 0.44b | 1.46 ± 0.49a |
| | Pt | 0.85 ± 0.17b | 1.25 ± 0.21b | 1.40 ± 0.41bc | 1.57 ± 0.50b | 0.83 ± 0.22b | 1.17 ± 0.32a | 1.85 ± 0.28a | 1.83 ± 0.23a |
| | Ct | 1.00 ± 0.26b | 1.99 ± 0.65a | 2.64 ± 0.67a | 2.25 ± 1.20ab | 0.80 ± 0.24b | 1.06 ± 0.32a | 1.21 ± 0.21b | 1.25 ± 0.48a |
| CAT | CjSz | 0.98 ± 0.20ab | 0.77 ± 0.10a | 1.14 ± 0.14ab | 1.00 ± 0.10ab | 0.91 ± 0.30a | 0.67 ± 0.09c | 0.87 ± 0.14a | 0.65 ± 0.10c |
| | CjZy | 0.74 ± 0.03b | 0.91 ± 0.09a | 1.52 ± 0.28a | 0.93 ± 0.08b | 0.57 ± 0.17a | 1.08 ± 0.17b | 0.71 ± 0.09ab | 1.56 ± 0.24a |
| | Cp | 0.90 ± 0.06ab | 0.79 ± 0.18a | 0.91 ± 0.04b | 1.08 ± 0.14ab | 0.70 ± 0.14a | 1.59 ± 0.10a | 0.90 ± 0.08a | 0.61 ± 0.03c |
| | Pt | 1.08 ± 0.11a | 0.72 ± 0.02a | 0.47 ± 0.06c | 1.28 ± 0.08a | 1.04 ± 0.14a | 0.84 ± 0.15bc | 0.53 ± 0.14b | 0.85 ± 0.10bc |
| | Ct | 0.77 ± 0.10 ab | 0.66 ± 0.12 a | 1.33 ± 0.44 a | 1.15 ± 0.15 ab | 0.79 ± 0.11 a | 0.88 ± 0.11 bc | 0.70 ± 0.13 ab | 1.09 ± 0.19 b |

Note: Data showed in the table are expressed as ratios relative to the values obtained on control seedling. Different lowercase letters indicate significant differences at $p < 0.05$.

Table 3. Mean comparison of soluble proteins and soluble sugars.

| Species | Flooding stress | | Drought stress | | Alkaline stress | | Freezing stress | | |
|------------------|-----------------|---------------|----------------|---------------|-----------------|---------------|-----------------|---------------|---------------|
| | 6 weeks | 7 weeks | 1 week | 2 weeks | pH = 8.0 | pH = 9.0 | 1 h | 2 h | |
| Soluble proteins | CjSz | 1.11 ± 0.17bc | 1.10 ± 0.26d | 1.35 ± 0.36a | 1.52 ± 0.08ab | 1.86 ± 0.18a | 2.04 ± 0.17a | 2.15 ± 0.18a | 2.16 ± 0.17a |
| | CjZy | 1.01 ± 0.17c | 1.46 ± 0.28bc | 1.19 ± 0.19a | 1.17 ± 0.28b | 1.81 ± 0.06a | 1.82 ± 0.31a | 1.58 ± 0.09c | 1.77 ± 0.32b |
| | Cp | 1.59 ± 0.19a | 1.88 ± 0.19a | 1.29 ± 0.32a | 1.67 ± 0.28a | 1.62 ± 0.10b | 1.83 ± 0.18a | 1.74 ± 0.17bc | 1.77 ± 0.17b |
| | Pt | 1.09 ± 0.07bc | 1.24 ± 0.17cd | 1.13 ± 0.19a | 1.11 ± 0.17b | 1.13 ± 0.13c | 1.19 ± 0.26b | 1.84 ± 0.09b | 1.93 ± 0.10ab |
| | Ct | 1.26 ± 0.02b | 1.54 ± 0.12b | 1.21 ± 0.55a | 1.35 ± 0.51ab | 1.23 ± 0.15c | 1.30 ± 0.15b | 1.28 ± 0.23d | 1.36 ± 0.17c |
| Soluble sugars | CjSz | 2.78 ± 0.48a | 3.24 ± 0.40a | 2.50 ± 0.62a | 1.87 ± 0.71bc | 1.77 ± 0.23ab | 2.08 ± 0.51b | 1.37 ± 0.20a | 1.25 ± 0.21b |
| | CjZy | 1.52 ± 0.45c | 1.79 ± 0.10b | 1.36 ± 0.37b | 1.62 ± 0.13bc | 1.42 ± 0.37bc | 2.32 ± 0.44ab | 1.29 ± 0.13a | 1.53 ± 0.23ab |
| | Cp | 2.80 ± 0.15a | 2.83 ± 0.53a | 1.96 ± 0.30ab | 2.76 ± 0.92a | 2.22 ± 0.63a | 2.77 ± 0.37a | 1.14 ± 0.41a | 1.74 ± 0.22a |
| | Pt | 2.12 ± 0.49b | 2.13 ± 0.33b | 1.86 ± 0.13ab | 1.26 ± 0.43c | 0.98 ± 0.16c | 1.15 ± 0.16c | 1.23 ± 0.28a | 1.41 ± 0.32ab |
| | Ct | 3.06 ± 0.26a | 3.16 ± 0.64a | 2.44 ± 0.93a | 2.25 ± 0.17ab | 1.01 ± 0.24c | 1.18 ± 0.25c | 1.17 ± 0.33a | 1.63 ± 0.31a |

Note: Data shown in the table are expressed as ratios relative to the values obtained on control seedling. Different lowercase letters indicate significant differences at $p < 0.05$.

parameters. By comparing the comprehensive evaluation values of all citrus rootstocks, we observed that the ranking of tolerance for each rootstock varied with treatment time. The comprehensive evaluation values were determined using a membership function and weight calculation, which allowed us to assess the performance of each rootstock under each different abiotic stress treatment (Table 4).

The results showed that CjSz had the highest comprehensive evaluation values during 6 and 7 weeks of flooding (0.710 and 0.966), at pH 8.0 (0.810), and after 2 h of freezing treatment (0.749). On the other hand, Ct exhibited the lowest comprehensive evaluation values during 6 weeks of flooding (0.141), and 2 weeks of drought (0.171), at pH 9.0 (0.272), and after 2 h of freezing treatment (0.018) (Table 4).

Performances of different graft combinations

Among the different graft combinations, the success rate of grafting 'Chunjian' and 'Mingrijian' onto Pt rootstocks was below 80%, with success rates of 70.67% and 79.33%, respectively (Table 5). The highest survival rates of 100% were

observed in 'Buzhihuo' was grafted onto CjSz and 'Dafen' grafted onto Ct. In terms of graft joint thickness (Ta), 'Dafen' grafted onto CjSz had the highest value of 20.13 mm, while 'Mingrijian' grafted onto Pt had the lowest value of 6.65 mm. The stem thickness below the graft joint (Tb) was the highest in 'Mingrijian' grafted onto CjSz (28.62 mm) and the lowest in 'Buzhihuo' grafted onto Pt (11.80 mm). The Ta/Tb ratio, which indicates the relative thickness above the blow the graft joint, was the highest in 'Tarocco' grafted onto CjZy and 'Tarocco' grafted onto Ct (0.82 for both), followed by 'Dafen' and 'Buzhihuo' grafted onto Ct (0.81 and 0.80, respectively). The lowest Ta/Tb ratio was observed in 'Mingrijian' and 'Chunjian' grafted onto Pt (0.52 and 0.58, respectively).

The tree growth was significantly affected by different rootstocks (Table 6). The citrus scions grafted onto CjSz rootstock exhibited strong tree vigor, followed by Ct, CjZy, and Pt (Table 6). The leaf greenness, as indicated by the SPAD value, was the highest in trees with 'Tarocco' grafted onto CjZy (86.46) and 'Buzhihuo' grafted onto CjSz (85.79), while it was lowest for 'Mingrijian', 'Chunjian', and 'Buzhihuo' grafted onto Pt (71.43,

Table 4. Comprehensive evaluation of five genotypes citrus rootstocks under different stresses.

| Treatment | Variety | Comprehensive evaluation | | | Membership function | | | Comprehensive evaluation value | Order |
|-------------------|---------|--------------------------|--------|--------|---------------------|-------|-------|--------------------------------|-------|
| | | F1 | F2 | F3 | U1 | U2 | U3 | | |
| Flooding 6 weeks | CjSz | 2.778 | -0.145 | 0.154 | 1.000 | 0.397 | 0.677 | 0.710 | 1 |
| | CjZy | -0.703 | -1.678 | 0.998 | 0.130 | 0.000 | 1.000 | 0.229 | 4 |
| | Cp | -1.222 | 0.620 | 0.569 | 0.000 | 0.596 | 0.836 | 0.375 | 3 |
| | Pt | -0.270 | 2.180 | -0.108 | 0.238 | 1.000 | 0.577 | 0.592 | 2 |
| | Ct | -0.582 | -0.978 | -1.614 | 0.160 | 0.181 | 0.000 | 0.141 | 5 |
| | Weights | | | | 0.440 | 0.388 | 0.172 | | |
| Flooding 7 weeks | CjSz | 2.261 | 1.481 | 0.119 | 1.000 | 1.000 | 0.621 | 0.966 | 1 |
| | CjZy | 0.985 | -1.106 | 0.866 | 0.715 | 0.000 | 1.000 | 0.565 | 2 |
| | Cp | -1.878 | -0.464 | 0.269 | 0.075 | 0.248 | 0.698 | 0.174 | 5 |
| | Pt | -2.215 | 1.047 | -0.146 | 0.000 | 0.832 | 0.487 | 0.249 | 4 |
| | Ct | 0.847 | -0.957 | -1.108 | 0.684 | 0.058 | 0.000 | 0.468 | 3 |
| | Weights | | | | 0.663 | 0.246 | 0.091 | | |
| Drought 1 week | CjSz | -0.468 | 1.581 | 0.303 | 0.239 | 1.000 | 0.504 | 0.516 | 2 |
| | CjZy | -1.545 | 0.762 | -0.946 | 0.000 | 0.755 | 0.000 | 0.228 | 4 |
| | Cp | 0.003 | -1.758 | -0.943 | 0.343 | 0.000 | 0.001 | 0.178 | 5 |
| | Pt | 2.964 | 0.349 | 0.050 | 1.000 | 0.631 | 0.401 | 0.781 | 1 |
| | Ct | -0.954 | -0.934 | 1.535 | 0.131 | 0.247 | 1.000 | 0.321 | 3 |
| | Weights | | | | 0.519 | 0.302 | 0.179 | | |
| Drought 2 weeks | CjSz | -0.005 | 1.386 | 1.251 | 0.386 | 1.000 | 1.000 | 0.672 | 2 |
| | CjZy | -1.787 | 0.898 | -1.024 | 0.000 | 0.864 | 0.000 | 0.287 | 4 |
| | Cp | 0.005 | 0.146 | -0.150 | 0.388 | 0.654 | 0.384 | 0.476 | 3 |
| | Pt | 2.826 | -0.236 | -0.492 | 1.000 | 0.547 | 0.234 | 0.747 | 1 |
| | Ct | -1.039 | -2.194 | 0.415 | 0.162 | 0.000 | 0.632 | 0.171 | 5 |
| | Weights | | | | 0.535 | 0.332 | 0.133 | | |
| Alkaline pH = 8.0 | CjSz | 0.492 | 1.831 | 0.873 | 0.674 | 1.000 | 1.000 | 0.810 | 1 |
| | CjZy | 2.056 | 0.273 | -0.941 | 1.000 | 0.551 | 0.000 | 0.761 | 2 |
| | Cp | 1.059 | -1.087 | 0.226 | 0.792 | 0.160 | 0.643 | 0.574 | 3 |
| | Pt | -2.749 | 0.626 | -0.598 | 0.000 | 0.653 | 0.189 | 0.229 | 5 |
| | Ct | -0.859 | -1.643 | 0.440 | 0.393 | 0.000 | 0.761 | 0.301 | 4 |
| | Weights | | | | 0.582 | 0.323 | 0.094 | | |
| Alkaline pH = 9.0 | CjSz | 0.178 | 1.765 | -0.992 | 0.681 | 1.000 | 0.000 | 0.699 | 2 |
| | CjZy | 1.244 | 0.969 | 0.689 | 0.941 | 0.770 | 0.990 | 0.885 | 1 |
| | Cp | 1.484 | -1.154 | 0.590 | 1.000 | 0.156 | 0.932 | 0.676 | 3 |
| | Pt | -2.614 | 0.115 | 0.706 | 0.000 | 0.523 | 1.000 | 0.342 | 4 |
| | Ct | -0.292 | -1.695 | -0.993 | 0.567 | 0.000 | 0.000 | 0.272 | 5 |
| | Weights | | | | 0.481 | 0.372 | 0.148 | | |
| Freezing 1 h | CjSz | -0.127 | 0.600 | 1.842 | 0.330 | 0.644 | 1.000 | 0.568 | 2 |
| | CjZy | -1.364 | -1.396 | -0.133 | 0.021 | 0.000 | 0.284 | 0.068 | 5 |
| | Cp | 2.549 | 0.457 | -0.447 | 1.000 | 0.598 | 0.170 | 0.701 | 1 |
| | Pt | -1.447 | 1.702 | -0.916 | 0.000 | 1.000 | 0.000 | 0.324 | 3 |
| | Ct | 0.389 | -1.364 | -0.346 | 0.459 | 0.010 | 0.207 | 0.263 | 4 |
| | Weights | | | | 0.473 | 0.324 | 0.203 | | |
| Freezing 2 h | CjSz | 1.279 | 0.579 | -0.031 | 0.907 | 0.650 | 0.416 | 0.749 | 1 |
| | CjZy | -1.946 | 1.708 | 0.603 | 0.000 | 1.000 | 0.714 | 0.443 | 4 |
| | Cp | 0.900 | -1.426 | 1.209 | 0.800 | 0.029 | 1.000 | 0.567 | 3 |
| | Pt | 1.609 | 0.661 | -0.915 | 1.000 | 0.676 | 0.000 | 0.746 | 2 |
| | Ct | -1.843 | -1.521 | -0.866 | 0.029 | 0.000 | 0.023 | 0.018 | 5 |
| | Weights | | | | 0.516 | 0.340 | 0.144 | | |

75.63, and 75.70, respectively). Overall, the results suggest that CjSz exhibited good graft compatibility with the test scions.

Discussion

Grafting is widely used in citrus propagation and provides many agronomical advantages to scion^[32,33]. Rootstock is vital for the citrus industry as it provides resistance to multiple stresses^[3,34]. The rootstocks used in citrus have certain issues with stress resistance, disease resistance, or grafting compatibility^[14-16], rendering them inflexible to the varying soil environment and climatic conditions, and consequently,

cannot be widely used in various cultivars^[17]. Therefore, citrus rootstock cultivation and evaluation are critical for the industry's healthy, stable, and sustainable development^[27].

In this study, abiotic stress altered the physiological, metabolic, and molecular processes^[17,35]. Almost all rootstock leaf photosynthetic pigments (*Chl a*, *Chl b*, total *Chl*, and carotenoid) were decreased under abiotic stress treatments (Table 1), creating an imbalance in the photosynthetic machinery^[36]. Under stress conditions, MDA, soluble protein, soluble sugar contents, and antioxidant enzyme activities were in an unbalanced equilibrium state^[23,24,37]. Alkaline stress

Table 5. Survival rate and graft union situation of different graft combinations.

| Graft combination | | Survival rate (%) | Diameter of scion (mm) | Diameter of rootstock (mm) | Ration of scion to rootstock |
|-------------------|------------|-------------------|------------------------|----------------------------|------------------------------|
| Rootstock | Scion | | | | |
| CjSz | Chunjian | 95.67 | 18.81 ± 0.35ab | 26.52 ± 2.75b | 0.70 ± 0.03ab |
| | Buzhihuo | 100.00 | 18.31 ± 2.34ab | 23.96 ± 2.49c | 0.77 ± 0.02a |
| | Mingrijian | 91.33 | 17.87 ± 1.95b | 28.62 ± 2.88a | 0.63 ± 0.06c |
| | Dafen | 90.67 | 20.13 ± 2.52a | 28.40 ± 3.42ab | 0.70 ± 0.06ab |
| | Tarocco | 95.50 | 20.10 ± 1.68a | 30.12 ± 2.74a | 0.67 ± 0.01bc |
| CjZy | Chunjian | 86.33 | 11.92 ± 0.94b | 16.00 ± 0.46b | 0.75 ± 0.05ab |
| | Buzhihuo | 95.50 | 12.59 ± 1.87b | 16.05 ± 1.39b | 0.78 ± 0.06ab |
| | Mingrijian | 81.67 | 11.26 ± 1.28b | 16.01 ± 1.52b | 0.71 ± 0.01b |
| | Dafen | 95.50 | 14.78 ± 1.99a | 20.48 ± 2.29a | 0.74 ± 0.12ab |
| | Tarocco | 90.67 | 14.82 ± 2.01a | 17.66 ± 2.85ab | 0.82 ± 0.05a |
| Ct | Chunjian | 91.33 | 15.94 ± 1.64ab | 21.17 ± 1.71a | 0.75 ± 0.03ab |
| | Buzhihuo | 95.67 | 14.77 ± 2.50b | 18.83 ± 2.51ac | 0.80 ± 0.12ab |
| | Mingrijian | 90.67 | 15.02 ± 2.48b | 20.91 ± 2.27ab | 0.72 ± 0.05b |
| | Dafen | 100.00 | 16.44 ± 0.81ab | 21.60 ± 3.82a | 0.81 ± 0.09a |
| | Tarocco | 87.33 | 17.37 ± 1.07a | 21.28 ± 1.65a | 0.82 ± 0.03a |
| Pt | Chunjian | 70.67 | 8.19 ± 0.49bc | 14.10 ± 2.02abc | 0.58 ± 0.05ab |
| | Buzhihuo | 95.67 | 6.66 ± 0.66c | 11.80 ± 0.44c | 0.57 ± 0.07ab |
| | Mingrijian | 79.33 | 6.65 ± 0.76c | 12.81 ± 1.13bc | 0.52 ± 0.06b |
| | Dafen | 91.67 | 10.94 ± 0.97a | 17.01 ± 1.67a | 0.65 ± 0.07a |
| | Tarocco | 91.33 | 10.36 ± 1.84ab | 16.55 ± 0.68ab | 0.62 ± 0.18a |

Table 6. The growth situation of different graft combinations.

| Graft combination | | Scion length (cm) | Crown breadth | | Leaf greenness (SPAD) |
|-------------------|------------|-------------------|-------------------|-----------------|-----------------------|
| Rootstock | Scion | | Longitudinal (cm) | Horizontal (cm) | |
| CjSz | Chunjian | 97.54 ± 9.53b | 87.83 ± 14.42a | 89.18 ± 16.32b | 82.98 ± 0.74abc |
| | Buzhihuo | 79.11 ± 10.63d | 73.67 ± 5.78b | 78.00 ± 17.04c | 85.79 ± 1.85a |
| | Mingrijian | 115.78 ± 8.88a | 91.33 ± 13.53a | 110.38 ± 19.01a | 84.97 ± 0.47a |
| | Dafen | 88.11 ± 5.98c | 97.33 ± 20.63a | 98.00 ± 18.89ab | 81.34 ± 0.14c |
| | Tarocco | 81.11 ± 12.07cd | 61.56 ± 11.59c | 65.56 ± 9.19d | 84.36 ± 1.55ab |
| CjZy | Chunjian | 69.67 ± 11.15ab | 66.99 ± 4.20a | 66.88 ± 15.99a | 82.16 ± 2.12b |
| | Buzhihuo | 55.89 ± 9.71c | 52.33 ± 3.71b | 61.11 ± 3.20ab | 82.93 ± 0.57b |
| | Mingrijian | 69.67 ± 5.93ab | 58.33 ± 14.25ab | 71.00 ± 13.91a | 79.54 ± 3.83c |
| | Dafen | 73.56 ± 6.00a | 72.17 ± 16.54a | 71.33 ± 18.34a | 84.93 ± 3.30ab |
| | Tarocco | 57.56 ± 10.84bc | 64.22 ± 11.52a | 52.67 ± 20.54b | 86.46 ± 0.80a |
| Ct | Chunjian | 96.11 ± 10.24a | 77.44 ± 12.30a | 83.33 ± 14.88ab | 81.37 ± 1.92ab |
| | Buzhihuo | 82.44 ± 15.61b | 62.56 ± 8.55bc | 69.22 ± 14.34b | 81.36 ± 1.68abc |
| | Mingrijian | 97.89 ± 9.34a | 76.89 ± 5.43ab | 77.67 ± 31.56ab | 78.81 ± 1.97bcd |
| | Dafen | 87.33 ± 5.46ab | 89.22 ± 1.26a | 92.67 ± 20.85a | 83.47 ± 1.70a |
| | Tarocco | 73.22 ± 13.77c | 55.22 ± 12.85c | 53.33 ± 16.68c | 83.43 ± 1.28a |
| Pt | Chunjian | 41.18 ± 7.44ab | 38.09 ± 12.96ab | 42.13 ± 14.27a | 75.63 ± 2.67b |
| | Buzhihuo | 25.56 ± 3.17c | 18.89 ± 1.92c | 15.56 ± 1.50bc | 75.70 ± 2.04b |
| | Mingrijian | 32.11 ± 7.93bc | 20.78 ± 5.23bc | 15.11 ± 2.12c | 71.43 ± 4.99c |
| | Dafen | 51.44 ± 4.17a | 50.56 ± 10.40a | 48.00 ± 6.66a | 80.92 ± 4.33a |
| | Tarocco | 42.13 ± 16.27ab | 27.89 ± 10.00bc | 33.89 ± 15.79ab | 81.71 ± 0.70a |

inhibits plant growth far more than salt stress^[38]. CjZy is widely used as an alkaline-tolerant citrus rootstock in calcareous soil areas in China^[5]. Under alkaline stress, CjSz performed similarly to CjZy. Low temperatures cause the leaves to wilt and dehydrate, reducing the photosynthesis rate^[39]. The disruption of photosynthetic mechanisms causes excessive production of ROS, leading to oxidative stress, one of the most damaging consequences of freezing stress. Pt has shown the highest resistance to cold stress^[40–42]. CjSz and Pt undergo similar physiological and biochemical changes, indicating that CjSz is also highly resistant to freezing. Flooding is a seasonal stress factor affecting Chinese citrus production areas^[2]. Under flooding stress, the MDA content in Ct was the highest, and that in Pt was the lowest. Comprehensive analysis revealed that Cp is the

most flood-resistant genotype, correlating with previous research findings^[43]. CjSz had reasonable flooding resistance. Cp has been considered drought-resistant citrus rootstocks owing to its higher chlorophyll content and POD activity under drought stress than CjSz. CjSz had less MDA content than Cp but more soluble sugar and soluble protein content. The comprehensive evaluation revealed that CjSz's drought tolerance was second only to Pt and superior to other rootstocks.

Rootstocks significantly affect tree performance in multiple aspects^[9]. Several studies have indicated that rootstocks have a significant effect on shoot growth^[3,18]. In this study, the effects of rootstocks on the horticultural performance of scion varieties were investigated. Rootstock genotypes influence compatibility^[9]. In our study, the graft success rate of CjSz with

five citrus cultivars ranged from 91.33% to 100%. Scions on CjSz developed faster than those on Pt, suggests that graft compatibility is related to the genetic relationship between scion and rootstock. The supply of root-derived nutrients, such as water and minerals, to the shoots may be limited due to incompatibility, leading to poor shoot growth and leaf function^[44]. Scions on Pt exhibited a much smaller canopy size and lower SPAD value than those on other rootstocks, consistent with the findings in three late-ripening navel oranges^[45] and Folha Murcha sweet oranges^[46]. These results suggest that Pt can be used for dense planting and that two genotypes of *C. junos* are preferable for sparse planting, consistent with previous findings^[45]. Therefore, scion-rootstock compatibility based on graft success and tree vigor supports that 'Shuzhen No.1' has a high potential for usage as a citrus rootstock.

Author contributions

The authors confirm contribution to the paper as follows: conceptualization and supervision: Wang X; methodology: He W; investigation: Chai J, Wang Y, Wu Z, Li M, Lin Y, Luo Y, Yong Zhang, Yunting Zhang, Wang H; bioinformatic analyses: He W, Chai J; data curation: He W, Xie R, Chai J; manuscript preparation: He W; writing—review and editing: He W, Tang H, Wang X. All authors reviewed the results and approved the final version of the manuscript.

Data availability

Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

Acknowledgments

This work was financially supported by Sichuan Provincial Postdoctoral Science Foundation, Sichuan Science and Technology Program (2020ZHCG0027), and Shuangzhi Project Innovation Team of Sichuan Agricultural University (Grant No. P202107).

Conflict of interest

The authors declare that they have no conflict of interest.

Dates

Received 3 September 2023; Accepted 13 November 2023; Published online 1 February 2024

References

1. Wu GA, Terol J, Ibanez V, López-García A, Pérez-Román E, et al. 2018. Genomics of the origin and evolution of *Citrus*. *Nature* 554:311–16
2. Xie R, Zheng L, Jiao Y, Huang X. 2021. Understanding physiological and molecular mechanisms of citrus rootstock seedlings in response to root zone hypoxia by RNA-Seq. *Environmental and Experimental Botany* 192:104647
3. Balfagón D, Rambla JL, Granell A, Arbona V, Gómez-Cadenas A. 2022. Grafting improves tolerance to combined drought and heat stresses by modifying metabolism in citrus scion. *Environmental and Experimental Botany* 195:104793
4. Primo-Capella A, Forner-Giner MÁ, Martínez-Cuenca M, Terol J. 2022. Comparative transcriptomic analyses of citrus cold-resistant vs. sensitive rootstocks might suggest a relevant role of ABA signaling in triggering cold scion adaption. *BMC Plant Biology* 22:209
5. Wu J, Cao J, Su M, Feng G, Xu Y, et al. 2019. Genome-wide comprehensive analysis of transcriptomes and small RNAs offers insights into the molecular mechanism of alkaline stress tolerance in a citrus rootstock. *Horticulture Research* 6:33
6. Fan Z, Wu Y, Zhao L, Fu L, Deng L, et al. 2022. MYB308-mediated transcriptional activation of plasma membrane H⁺-ATPase 6 promotes iron uptake in citrus. *Horticulture Research* 9:uhac88
7. Habibi F, Liu T, Folta K, Sarkhosh A. 2022. Physiological, biochemical, and molecular aspects of grafting in fruit trees. *Horticulture Research* 9:uhac32
8. Yang L, Xia L, Zeng Y, Han Q, Zhang S. 2022. Grafting enhances plants drought resistance: current understanding, mechanisms, and future perspectives. *Frontiers in Plant Science* 13:1015317
9. Warschefsky EJ, Klein LL, Frank MH, Chitwood DH, Londo JP, et al. 2016. Rootstocks: Diversity, domestication, and impacts on shoot phenotypes. *Trends in Plant Science* 21:418–37
10. Raiol-Junior LL, de Carvalho EV, Moreira AS, Marques JPR, Stuchi ES, et al. 2022. Graft compatibility classification within Aurantioideae based on biometric traits and the anatomy of graft union. *Agriculture* 12:76
11. Shimada T, Endo T, Fujii H, Nakano M, Sugiyama A, et al. 2018. MITE insertion-dependent expression of *CitRKD1* with a RWP-RK domain regulates somatic embryogenesis in citrus nucellar tissues. *BMC Plant Biology* 18:166
12. Wang X, Xu Y, Zhang S, Cao L, Huang Y, et al. 2017. Genomic analyses of primitive, wild and cultivated citrus provide insights into asexual reproduction. *Nature Genetics* 49:765–72
13. Peng Z, Bredeson JV, Wu GA, Shu S, Rawat N, et al. 2020. A chromosome-scale reference genome of trifoliolate orange (*Poncirus trifoliata*) provides insights into disease resistance, cold tolerance and genome evolution in *Citrus*. *The Plant Journal* 104:1215–32
14. He W, Xie R, Luo L, Chai J, Wang H, et al. 2022. Comparative transcriptomic analysis of inarching invigorating rootstock onto incompatible grafts in *Citrus*. *International Journal of Molecular Sciences* 23:14523
15. He W, Xie R, Wang Y, Chen Q, Wang H, et al. 2022. Comparative transcriptomic analysis on compatible/incompatible grafts in *Citrus*. *Horticulture Research* 9:uhab72
16. He W, Wang Y, Chen Q, Sun B, Tang H, et al. 2018. Dissection of the mechanism for compatible and incompatible graft combinations of *Citrus grandis* (L.) Osbeck ('Hongmian Miyou'). *International Journal of Molecular Sciences* 19:505
17. Zhu S, Nong J, Luo G, Li Q, Wang F, et al. 2021. Varied tolerance and different responses of five citrus rootstocks to acid stress by principle component analysis and orthogonal analysis. *Scientia Horticulturae* 278:109853
18. Balfagón D, Terán F, de Oliveira TDR, Santa-Catarina C, Gómez-Cadenas A. 2022. Citrus rootstocks modify scion antioxidant system under drought and heat stress combination. *Plant Cell Reports* 41:593–602
19. Zhu S, Chen J, Ma Y, Yan S, Zhong G. 2013. Advances in the studies on citrus rootstock evaluation and application. *Acta Horticulturae Sinica* 40:1669–78
20. Jiang J, Hou R, Yang N, Li L, Deng J, et al. 2021. Physiological and TMT-labeled proteomic analyses reveal important roles of sugar and secondary metabolism in *Citrus junos* under cold stress. *Journal of Proteomics* 237:104145
21. Hasanuzzaman M, Nahar K, Alam MM, Roychowdhury R, Fujita M. 2013. Physiological, biochemical, and molecular mechanisms of heat stress tolerance in plants. *International Journal of Molecular Sciences* 14:9643–84

Evaluation of *Citrus junos* 'Shuzhen No.1'

22. Dumanović J, Nepovimova E, Natić M, Kuča K, Jačević V. 2021. The significance of reactive oxygen species and antioxidant defense system in plants: a concise overview. *Frontiers in Plant Science* 11:552969
23. Gill SS, Tuteja N. 2010. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiology and Biochemistry* 48:909–30
24. Ozgur R, Uzilday B, Sekmen AH, Turkan I. 2013. Reactive oxygen species regulation and antioxidant defence in halophytes. *Functional Plant Biology* 40:832–47
25. Jiang M, Zhang J. 2002. Water stress - induced abscisic acid accumulation triggers the increased generation of reactive oxygen species and up - regulates the activities of antioxidant enzymes in maize leaves. *Journal of Experimental Botany* 53:2401–10
26. Xie R, He W, Chai J, Luo L, Wang Y, et al. 2022. A study of scion phenotypes in pummelo grafted onto a new citrus rootstock *Citrus junos* 'Pujiang Xiangcheng'. *Horticulturae* 8:1039
27. Fu X, Huang X, Chen T, Zhang J, Wang Y, et al. 2017. A new citrus rootstock 'Pujiang Xiangcheng' (*Citrus junos*). *Journal of Fruit Science* 34:917–20
28. Wellburn AR, Lichtenthaler H. 1984. Formulae and program to determine total carotenoids and chlorophylls a and b of leaf extracts in different solvents. In *Advances in Photosynthesis Research. Advances in Agricultural Biotechnology*, ed. Sybesma C, vol 2. Springer, Dordrecht. pp. 9–12. https://doi.org/10.1007/978-94-017-6368-4_3
29. Shahid MA, Balal RM, Khan N, Simón-Grao S, Alfosea-Simón M, et al. 2019. Rootstocks influence the salt tolerance of Kinnow mandarin trees by altering the antioxidant defense system, osmolyte concentration, and toxic ion accumulation. *Scientia Horticulturae* 250:1–11
30. Landi M. 2017. Commentary to: "Improving the thiobarbituric acid-reactive-substances assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other interfering compounds" by Hodges et al., *Planta* (1999) 207:604–611. *Planta* 245:1067
31. Zaher-Ara T, Boroomand N, Sadat-Hosseini M. 2016. Physiological and morphological response to drought stress in seedlings of ten citrus. *Trees* 30:985–93
32. Ko D, Helariutta Y. 2017. Shoot–root communication in flowering plants. *Current Biology* 27:R973–R978
33. Dong D, Shi Y, Mou Z, Chen S, Zhao D. 2022. Grafting: a potential method to reveal the differential accumulation mechanism of secondary metabolites. *Horticulture Research* 9:uhac50
34. Mahmoud LM, Huyck PJ, Vincent CI, Gmitter FG Jr, Grosser JW, et al. 2021. Physiological responses and gene expression patterns in open-pollinated seedlings of a pummelo-mandarin hybrid rootstock exposed to salt stress and Huanglongbing. *Plants* 10:1439
35. Ziogas V, Tanou G, Filippou P, Diamantidis G, Vasilakakis M, et al. 2013. Nitrosative responses in citrus plants exposed to six abiotic stress conditions. *Plant Physiology and Biochemistry* 68:118–26
36. Lawson T, Blatt MR. 2014. Stomatal size, speed, and responsiveness impact on photosynthesis and water use efficiency. *Plant Physiology* 164:1556–70
37. García-Caparrós P, De Filippis L, Gul A, Hasanuzzaman M, Ozturk M, et al. 2021. Oxidative stress and antioxidant metabolism under adverse environmental conditions: a review. *The Botanical Review* 87:421–66
38. Yang C, Wang P, Li C, Shi D, Wang D. 2008. Comparison of effects of salt and alkali stresses on the growth and photosynthesis of wheat. *Photosynthetica* 46:107–14
39. Pietrini F, Chaudhuri D, Thapliyal AP, Massacci A. 2005. Analysis of chlorophyll fluorescence transients in mandarin leaves during a photo-oxidative cold shock and recovery. *Agriculture, Ecosystems & Environment* 106:189–98
40. Khan M, Hu J, Dahren B, Ming R, Zhang Y, et al. 2021. ERF108 from *Poncirus trifoliata* (L.) Raf. functions in cold tolerance by modulating raffinose synthesis through transcriptional regulation of *PtrRafS*. *The Plant Journal* 108:705–24
41. Zhang Y, Ming R, Khan M, Wang Y, Dahren B, et al. 2022. ERF9 of *Poncirus trifoliata* (L.) Raf. undergoes feedback regulation by ethylene and modulates cold tolerance via regulating a *glutathione S-transferase U17* gene. *Plant Biotechnology Journal* 20:183–200
42. Geng J, Wei T, Wang Y, Huang X, Liu J, et al. 2019. Overexpression of *PtrbHLH*, a basic helix-loop-helix transcription factor from *Poncirus trifoliata*, confers enhanced cold tolerance in pummelo (*Citrus grandis*) by modulation of H₂O₂ level via regulating a *CAT* gene. *Tree Physiology* 39:2045–54
43. Arbona V, Hossain Z, López-Climent MF, Pérez-Clemente RM, Gómez-Cadenas A. 2008. Antioxidant enzymatic activity is linked to waterlogging stress tolerance in citrus. *Physiologia Plantarum* 132:452–66
44. De Carvalho DU, Leite RP Junior, Yada IFU, Tazima ZH. 2022. Trifoliolate orange-related rootstocks enhance the horticultural performance of 'Shamouti' sweet orange under humid subtropical condition. *Agriculture* 12:1782
45. Zhu S, Huang T, Yu X, Hong Q, Xiang J, et al. 2020. The effects of rootstocks on performances of three late-ripening navel orange varieties. *Journal of Integrative Agriculture* 19:1802–12
46. Cantuarias-Avilés T, Mourão Filho FDAA, Stuchi ES, Da Silva SR, Espinoza-Nuñez E. 2011. Horticultural performance of 'Folha Murcha' sweet orange onto twelve rootstocks. *Scientia Horticulturae* 129:259–65



Copyright: © 2024 by the author(s). Published by Maximum Academic Press, Fayetteville, GA. This article is an open access article distributed under Creative Commons Attribution License (CC BY 4.0), visit <https://creativecommons.org/licenses/by/4.0/>.