

Article

SlWRKY80-mediated jasmonic acid pathway positively regulates tomato resistance to saline–alkali stress by enhancing spermidine content and stabilizing Na⁺/K⁺ homeostasis

Chunyu Shang^{1,2,3,†}, Xiaoyan Liu^{1,2,3,†}, Guo Chen^{1,2,3}, Hao Zheng^{1,2,3}, Abid Khan⁴, Guobin Li^{1,2,3,*} and Xiaohui Hu^{1,2,3,*}¹College of Horticulture, Northwest A&F University, Yangling, Shaanxi, 712100, China²Key Laboratory of Protected Horticultural Engineering in Northwest, Ministry of Agriculture and Rural Affairs, Yangling, Shaanxi, 712100, China³Shaanxi Protected Agriculture Engineering Technology Research Centre, Yangling, Shaanxi, 712100, China⁴Department of Horticulture, The University of Haripur, Haripur, 22620, Pakistan*Corresponding authors. E-mail: liguobin@nwafu.edu.cn; hxx1977@nwafu.edu.cn

†These authors contributed equally to this work.

Abstract

Saline–alkali is an important abiotic stressor influencing tomato production. Exogenous methyl jasmonate (MeJA) is well known to increase tomato resistance to a variety of stresses, although its exact mechanism is yet unknown. In this study we confirmed that 22.5 $\mu\text{mol/l}$ MeJA could significantly improve the saline–alkali stress resistance of tomato. Saline–alkali (300 mM) stress increased the endogenous MeJA and jasmonic acid (JA) contents of tomato by 18.8 and 13.4%, respectively. Exogenous application of 22.5 $\mu\text{mol/l}$ MeJA increased the endogenous MeJA and JA contents in tomato by 15.2 and 15.9%, respectively. Furthermore, we found an important transcription factor, SlWRKY80, which responded to MeJA, and constructed its overexpressing and knockout lines through genetic transformation. It was found that SlWRKY80 actively regulated tomato resistance to saline–alkali stress, and the spraying of exogenous MeJA (22.5 $\mu\text{mol/l}$) reduced the sensitivity of SlWRKY80 knockout lines to saline–alkali stress. The SlWRKY80 protein directly combines with the promoter of SlSPDS2 and SlNHX4 to positively regulate the transcription of SlSPDS2 and SlNHX4, thereby promoting the synthesis of spermidine and Na⁺/K⁺ homeostasis, actively regulating saline–alkali stress. The augmentation of JA content led to a notable reduction of 70.6% in the expression of SlJAZ1, and the release of the SlWRKY80 protein interacting with SlJAZ1. In conclusion, we revealed the mechanism of exogenous MeJA in tomato stress resistance through multiple metabolic pathways, elucidated that exogenous MeJA further promotes spermidine synthesis and Na⁺/K⁺ homeostasis by activating the expression of SlWRKY80, which provides a new theoretical basis for the study of the JA stress resistance mechanism and the production of tomato.

Introduction

Around the world, the tomato (*Solanum lycopersicum* L.) is the most widely grown and consumed horticultural crop. In addition to moderate salt sensitivity, tomatoes are also susceptible to abiotic stresses such as saline–alkali, which negatively affect their growth [41]. However, there is a problem of salinization on 33% of the world's arable land, a condition that significantly impedes tomato productivity, rendering it a primary environmental concern thwarting global agricultural development of a high-quality nature [11]. Therefore, studying how tomato plants respond to saline–alkali stress is essential to improving tomato quality and yield.

Plants activate various mechanisms and trigger changes in endogenous phytohormones to response to saline–alkali stress, including jasmonic acid (JA) [74], abscisic acid [13, 39, 66, 74], brassinosteroids edna [32], and ethylene [35] etc. Plants under salt stress exhibit positive effects on JA, and endogenous JA content is

enhanced and JA signaling is activated under salt stress [69]. The high JA-accumulating tomato mutant *res* exhibits stronger salt tolerance [16], while *def-1* (JA-deficient mutant) was salt-sensitive [2]. Similarly, studies on JA-related mutants in wheat [75], rice [17], and corn (maize) [3] have shown that JA is associated with salt stress responses. Methyl jasmonate (MeJA) exhibits a similar function to JA in participating in plant stress resistance. MeJA enters the plant through the stomata, is hydrolyzed into JA by esterases in the cytoplasm, and facilitates long-distance signal propagation and interplant communication. This process induces defense responses in nearby plants, fortifying their resilience as well [60]. For instance, exogenously applied MeJA enhances plants' salt tolerance, either by maintaining reactive oxygen species (ROS) homeostasis, or by stabilizing the ion equilibrium [69]. The proteins of JAZ, which contain a jasmonate ZIM domain, act as repressors that participate in multiple signaling pathways and can bind to transcription factors or other corepressor proteins,

Received: 25 October 2023; Accepted: 16 January 2024; Published: 24 January 2024; Corrected and Typeset: 27 March 2024

© The Author(s) 2024. Published by Oxford University Press on behalf of Nanjing Agricultural University. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

linking the JA signaling pathway with other signaling pathways. CO11, an F-box protein, is a core component of the JA signaling receptor [5, 36].

Hormones exert direct effects on transcription factors, including the WRKY gene family, one of the first and largest transcriptional regulators to be identified. Furthermore, WRKY can mediate the influence of JAZ genes on *Arabidopsis thaliana*'s resistance to *Botrytis cinerea* [23]. Additionally, CaWRKY40 in chili pepper suppresses the expression of the JA signaling repressor JAZ8, thereby enhancing disease resistance [48]. As well as being involved in stress response, the WRKY proteins bind to the W-box (TTGAC-T/C) sequence in the promoter regions of target genes [49]. Factors including AtWRKY25/33 [25], AtWRKY8 [21], AtWRKY46 [14], MdWRKY100 [40], and AcWRKY28 [62] along with negative regulators such as AtWRKY15 [55], PalWRKY77 [26], ZmWRKY20/115 [10], and OsWRKY53 [70], are reported to be involved in salt stress responses. Notably, according to a recent study, WRKY transcription factors and their feedback loops function as central nodes in salt-responsive gene regulatory networks, indicating that WRKYs play an indispensable role in plant responses to salt stress [61].

There are 83 known SlWRKY genes in tomato [22]. A significant role is played by SlWRKY80 in the plant's disease resistance, answering the call of signals from salicylic acid (SA) as well as JA [44]. Group III of the SlWRKY family genes actively participate in the response to abiotic stress [7], and the group III subfamily of the tomato SlWRKY family encompasses a collective sum of eight genes, specifically identified as SlWRKY30, SlWRKY41, SlWRKY52, SlWRKY53, SlWRKY54, SlWRKY59, SlWRKY80, and SlWRKY81 [12]. The interaction of SlWRKY30 and SlWRKY80 further bolsters the resistance of SlPR-STH2 to bacterial wilt [12]. Yet the biological function of SlWRKY80 under abiotic threats such as saline-alkali stress remains unelucidated, and the stress response mechanisms of SlWRKY80 are not fully understood. Other genes in the WRKY family such as SlWRKY33 [76], SlWRKY39 [53], SlWRKY8 [15], SlWRKY79 [19], and SlWRKY23 [52] are associated with salt stress, while SlWRKY28 [57] has been associated with alkaline stress. However, current research on the SlWRKY genes in tomato under saline-alkali stress still needs further clarification. Shedding light on the pivotal function of WRKY genes under such adversities, and comprehending their operational mechanisms, will offer theoretical foundations for the enhancement of tomato resistance breeding, and has crucial significance.

Spermidine (Spd) is a free compound existing within plants, a type of polyamine, and has a prominent role in preventing ionic toxicity and reducing salt-alkali stress. Overexpression of SlSPDS2, which is involved in Spd synthesis, reduces Na⁺/K⁺ and H₂O₂ levels, mitigated ionic toxicity of tomato [56]. In response to abiotic stresses like cold, freezing, and salinity, Spd synthesis-related genes are overexpressed, along with transcription factors like WRKY that are increased [27]. Increasing the expression of PtSPD also significantly improved the tolerance of a member of the *Populus* genus, *P. davidiana*, to saline-alkali stress [57]. All these data indicate that the genes for WRKY TF and SPDS gene play important roles in plant resistance to saline-alkali stress, but the mechanism of action between WRKY TF and SPDS gene is still unclear.

Additionally, plants express genes involved in ion transport in the plasma membrane to combat saline-alkali stress, such as SOS1, HKT1.1, HKT1.2, NHX1, and NHX4 [8, 45]. These functional genes expel excessive sodium ions from the cells or sequester them into vacuoles, reducing sodium ion accumulation within the cells. Concurrently, the expression of potassium channel

protein-encoding genes LKT1, HAK20, and NHX2 is induced, which in turn promotes potassium ion absorption and transport. By doing so, the saline-alkali stress-induced ionic toxicity is lessened and the Na⁺/K⁺ ratio is decreased [18]. NHXs are membrane-localized proteins that play roles in maintaining the Na⁺/K⁺ and pH homeostasis within cells. A reduction in the concentration of Na⁺ in the cytosol is achieved by removing or enclosing Na⁺ ions from the cytoplasm. By activating potassium ion channel proteins and increasing K⁺ content, the NHX proteins preserve Na⁺/K⁺ ion homeostasis, constituting an essential mechanism for mitigating ionic toxicity and enhancing saline-alkali stress resistance [8]. Furthermore, AtWRKY75 is able to bind to the promoter of AtSOS1 in *Arabidopsis*, thereby regulating the expression of AtSOS1 [38].

We observed that the SlWRKY gene family of tomato showed significant responses to saline-alkali stress, and a certain concentration of exogenous MeJA could enhance tomato saline-alkali tolerance. Keeping in view the importance of the SlWRKY gene family, we selected the SlWRKY80 gene through transcriptome analysis. SlWRKY80-overexpressing and knockout lines were obtained through genetic transformation, and functional validation was conducted under saline-alkali stress. This experimental study found that the transcription factor SlWRKY80 was significantly upregulated under saline-alkali stress, and the promoter of SlWRKY80 can respond to both saline-alkali and exogenous MeJA signals simultaneously. This study aimed to investigate how exogenous MeJA participates in tomato saline-alkali tolerance through the regulation of SlWRKY80, and in order to find a theoretical explanation for tomato tolerance to saline-alkali stress we studied the relationship between SlWRKY80, JA signal transduction, Spd synthesis, and the balance of Na⁺/K⁺ homeostasis.

Results

Exogenous methyl jasmonate has dual effects on tomato saline-alkali stress

Several concentrations of MeJA were applied to wild-type (WT) tomato seedlings to investigate their response to saline-alkali stress. With higher concentrations of exogenous MeJA under saline-alkali stress, the tolerance of tomato seedlings initially exhibited an increasing trend followed by a decrease (Fig. 1A). Further analysis showed that when the concentration of exogenous MeJA was 22.5 μmol/l, the morphological indicators of stem diameter and plant height (Supplementary Data Fig. S1) and physiological indicators such as SOD and POD (Fig. 1B–F, Supplementary Data Fig. S2) were significantly higher than in other treatment groups, while the change trend of malondialdehyde content was opposite (Fig. 1G). Therefore, exogenous MeJA has dual effects on tomato saline-alkali stress, while spraying exogenous 22.5 μmol/l MeJA can significantly affect tomato resistance to saline-alkali stress.

SlWRKY80 responded to saline-alkali stress and methyl jasmonate

In tomato, SlWRKY80 significantly responded to saline-alkali stress when eight genes in the third subfamily of the SlWRKY family were examined (Fig. 2A) [12]. We further examined the relative expression level of SlWRKY80 in tomato seedlings treated with saline-alkali at 3, 6, 12, and 24 h. In the presence of saline-alkali treatment (S), SlWRKY80 expression levels were significantly increased. When treated with saline-alkali and exogenous MeJA (S+M), a significant increase in SlWRKY80 expression was observed (Fig. 2B).




Fig. 1. Impact of different concentrations of exogenous MeJA on tomato's resilience to saline-alkali stress. **A** Phenotypes of tomato plants sprayed with different concentrations of exogenous MeJA under saline-alkali treatment. **B** Sound seedling index. **C–E** SOD, POD, and CAT enzyme activities. **F** Proline content. **G** Malondialdehyde content. **H** Root activity. According to the LSD test, significant differences are indicated by lowercase letters ($P < 0.05$), and values indicate the mean across three biological replicates.

To verify whether the *SlWRKY80* promoter is regulated by saline-alkali and MeJA, we subjected pCambia1391-pro-*SlWRKY80*-positive lines to 300 mM saline-alkali treatment (+S), exogenous spraying of 22.5 $\mu\text{mol/l}$ MeJA (+MeJA), and simultaneous treatment with 300 mM saline-alkali and exogenous spraying of 22.5 $\mu\text{mol/l}$ MeJA (+S+MeJA). We found that the depth of GUS staining gradually deepened (Fig. 2C). Grayscale analysis of GUS-stained images yielded the same results (Supplementary Data Fig. S3). The data also showed that S+M treatment made the GUS staining of pCambia1391-pro-*SlWRKY80* transgenic material significantly higher than that of other treatments (Supplementary Data Fig. S3), indicating that the promoter of *SlWRKY80* responded to both saline-alkali treatment and exogenous MeJA, and S+MeJA treatment could make the promoter of *SlWRKY80* respond more significantly. In the analysis of the 2000-bp promoter upstream of *SlWRKY80*, we also found that there were four cis-acting elements responding to MeJA (Supplementary Data Fig. S4), which was also consistent with the GUS staining results in this experiment (Fig. 2C). *SlWRKY80* protein and GFP were fused under the PBI121 vector in order to determine its subcellular localization. A fluorescence microscope analysis revealed that GFP fluorescence was only found in the nucleus (Fig. 2D), indicating that *SlWRKY80* protein belongs to the nucleus.

SlWRKY80 positively regulates saline-alkali stress

SlWRKY80 was further tested under saline-alkali stress. We obtained *SlWRKY80*-overexpressing lines 80OE-1 and 80OE-3 (Supplementary Data Fig. S5) and knockout lines 80CR-3 and 80CR-4 (Supplementary Data Fig. S6) by genetic transformation.

Before saline-alkali stress, the lines did not differ significantly in plant height (Supplementary Data Fig. S7A), soluble sugar (Supplementary Data Fig. S8A), and soluble protein content (Supplementary Data Fig. S8B). However, stem diameter (Supplementary Data Fig. S7B) and leaf area (Supplementary Data Fig. S7C) of the *SlWRKY80*-overexpressing lines showed no significant differences. Carotenoids (Supplementary Data Fig. S8C), chlorophyll a (Supplementary Data Fig. S8D), and total chlorophyll content (Supplementary Data Fig. S8F) were significantly higher than those of WT, while the *SlWRKY80*-knockout line showed the opposite trend.

The *SlWRKY80* transgenic and WT tomato seedlings were treated with 300 mM saline-alkali solution, and obvious phenotypes were observed on the eighth day. Compared with WT, *SlWRKY80*-overexpressing lines showed significantly better growth, and the growth of *SlWRKY80* knockout lines was the worst (Fig. 3A). The sound seedling index (Fig. 3B), DAB staining (Fig. 3C), and NBT staining (Fig. 3D) also showed the same results. In




Fig. 2. Gene expression analysis of the third subgroup of SlWRKYs in tomato under saline-alkali stress, response of SlWRKY80 to saline-alkali stress and MeJA, and the subcellular localization of SlWRKY80. **A** Heat map of gene expression in the third subgroup of tomato SlWRKYs under saline-alkali stress. **B** SlWRKY80 expression under S, M and S + M conditions. S represents 300 mM saline-alkali stress; M represents exogenous spraying of 22.5 $\mu\text{mol/l}$ MeJA; S + M represents saline-alkali stress and exogenous spraying of 22.5 $\mu\text{mol/l}$ MeJA. According to the LSD test, significant differences are indicated by lowercase letters ($P < 0.05$), and values indicate the mean across three biological replicates. **C** GUS staining map of pCambia1391-pro-SlWRKY80 transgenic material under saline-alkali, exogenous MeJA spraying, and simultaneous treatment with saline-alkali and exogenous MeJA. S, saline-alkali; MeJA concentration was 22.5 $\mu\text{mol/l}$. **D** Subcellular localization of SlWRKY80 protein.

contrast to the WT, except for the lower content of malondialdehyde in the SlWRKY80-overexpressing line, its morphological and physiological characteristics were significantly higher, while the SlWRKY80-knockout line showed the opposite results, indicating that overexpression of SlWRKY80 significantly improved stem diameter, leaf area, antioxidant capacity, root activity, etc. of the tomato plants. Saline-alkali stress stimulated SlWRKY80 expression (Fig. 3E–J). Therefore, we concluded that SlWRKY80 can actively regulate saline-alkali stress.

Exogenous application of methyl jasmonate reduces the sensitivity of SlWRKY80 knockout lines to saline-alkali stress

There is a sensitivity to saline-alkali stress in SlWRKY80 knockout lines, and exogenous 22.5 $\mu\text{mol/l}$ MeJA significantly enhances the tomato's resistance to saline-alkali stress. We asked whether exogenous 22.5 $\mu\text{mol/l}$ MeJA reduces the SlWRKY80-knockout lines' sensitivity to saline-alkali stress.

Following 8 days of saline-alkali treatment, exogenous 22.5 $\mu\text{mol/l}$ MeJA significantly reduced the sensitivity of SlWRKY80 knockout lines to saline-alkali stress. Saline-alkali stress affected 80CR-3 lines the most compared with WT, and the damage of the 80CR-3 lines was alleviated by saline-alkali stress after spraying exogenous 22.5 $\mu\text{mol/l}$ MeJA (80CR-3 + MeJA) (Fig. 4A). Before treatment, SOD activity (Fig. 4D), POD activity (Fig. 4E), and CAT activity (Fig. 4F) did not differ significantly. Sound seedling index and SOD, POD, and CAT activities of tomato seedlings in the 80CR-3 + MeJA group were significantly higher than in the 80CR-3 group on the eighth day after treatment, while DAB staining (Fig. 4Ca) and NBT staining (Fig. 4Cb) showed the same results. These results showed that exogenous 22.5 $\mu\text{mol/l}$ MeJA could significantly reduce the sensitivity of SlWRKY80 knockout lines to saline-alkali stress.

SlWRKY80 participates in the stress resistance pathway of jasmonic acid to saline-alkali stress through its interaction with SlJAZ1

A study was conducted to determine if endogenous JA and MeJA are impacted by saline-alkali stress in SlWRKY80 transgenic lines. We found that SlWRKY80-overexpressing lines had significantly increased contents of endogenous MeJA and JA in tomato. After 24 h of saline-alkali treatment, SlWRKY80-overexpressing lines contained significantly more MeJA and JA than WT, while the SlWRKY80 knockout lines were the opposite (Fig. 5A). Therefore, SlWRKY80-overexpressing lines had significantly increased endogenous MeJA and JA levels, while the knockout lines showed the opposite effect. MeJA and JA levels were lowest in the 80CR-3 group, followed by 80CR-3 + MeJA, and highest in the WT group of tomatoes treated with exogenous 22.5 $\mu\text{mol/l}$ MeJA and saline-alkali (Fig. 5B). The 300 mM saline-alkali stress increased the endogenous MeJA and JA contents in tomato by 18.8 and 13.4%, respectively. Exogenous application of 22.5 $\mu\text{mol/l}$ MeJA increased the endogenous MeJA and JA contents in tomato by 15.2 and 15.9%, respectively. In addition, the findings also revealed a significant increase in the expression of JA synthesis-related genes, specifically SlLxD and SlAOC, upon exogenous MeJA spraying (Supplementary Data Fig. S11). Consequently, it can be inferred that exogenous application of 22.5 $\mu\text{mol/l}$ MeJA could significantly increase the content of endogenous JA.

JA content is negatively correlated with the expression of SlJAZs [30, 50]. Therefore, we have focused on the transcriptional suppressor genes SlJAZs in tomato, which are also significant in signal transduction. We verified the protein level interaction between SlWRKY80 and SlJAZ1, SlJAZ2 and SlJAZ5 through yeast two-hybrid (Y2H) assays (Fig. 5C). We verified it by luciferase complementary imaging (LCI) (Fig. 5D), bimolecular fluorescence complementation (BiFC) (Fig. 5E), and pull-down (Fig. 5F) assays




Fig. 3. Morphology and physiological responses of WT and SIWRKY80 transgenic plants to saline-alkali stress. **A** Phenotypic image of SIWRKY80 transgenic tomato lines treated with saline-alkali stress. Take photos of tomato seedlings after 8 days of saline-alkali treatment. **B** Sound seedling index. **C, D** DAB and NBT holding charts on the eighth day of saline-alkali stress. **E–G** SOD, POD, and CAT activity. **H–J** Proline content, malondialdehyde content, and root activity. According to the LSD test, significant differences are indicated by lowercase letters ($P < 0.05$), and values indicate the mean across three biological replicates.

in vivo and *in vitro*, and found that only SIJAZ1 could get positive results under verification by these molecular means. Therefore, we conclude that SIWRKY80 is involved in the resistance pathway of JA to saline-alkali stress through the interaction with SIJAZ1.

SIWRKY80 binds the promoter of SISPD2 to promote spermidine synthesis

To further investigate how saline-alkaline stress regulates SIWRKY80, we measured the content of endogenous Spd of tomato seedlings of different lines pre- and post-saline-alkali. There was significantly reduced endogenous Spd content in SIWRKY80 knockout lines on the eighth day. At the same time, overexpression of SIWRKY80 before saline-alkali treatment could significantly increase the content of tomato Spd, which was further enhanced

after 8 days of saline-alkali treatment (Fig. 6A). In addition, the expression trend of SISPD2 was similar to that of endogenous Spd content (Supplementary Data Fig. S9).

To explore whether this enhancement is regulated by SIWRKY80, we analyzed the 3000-bp promoter upstream of SISPD2, a functional gene for Spd synthesis, and found that it has three W-boxes. Subsequently, we confirmed that SIWRKY80 could directly bind to the three W-boxes in the SISPD2 promoter by *in vivo* and *in vitro* verification methods such as the yeast one hybrid (Y1H) assay (Fig. 6B), chromatin immunoprecipitation (ChIP)-qPCR (Fig. 6C), and the electrophoretic mobility shift assay (EMSA) (Fig. 6D). To sum up, tomato can enhance resistance to saline-alkali stress by enhancing the expression of SIWRKY80 and regulating the synthesis of Spd.




Fig. 4. Exogenous application of MeJA reduces the sensitivity of *SlWRKY80* knockout lines to saline-alkali stress. **A** Phenotypic map of the effect of exogenous MeJA on the *SlWRKY80* knockout lines on the eighth day after treatment. **B** Sound seedling index of WT, 80CR-3, and 80CR-3+MeJA. **C** DAB (a) and NBT (b) staining of WT, 80CR-3, 80CR-3+MeJA on the eighth day of saline-alkali treatment. **(D-F)** SOD, POD, and CAT activities of WT, 80CR-3, 80CR-3 + Break. According to the LSD test, significant differences are indicated by lowercase letters ($P < 0.05$), and values indicate the mean across three biological replicates.

SlWRKY80 binds to the *SINHX4* promoter and reduces Na^+/K^+ ratio

In order to further investigate whether *SlWRKY80* regulates saline-alkali stress by regulating Na^+ and K^+ transport, we found through experiments that after 8 days of saline-alkali stress treatment the Na^+ content, K^+ content, and Na^+/K^+ in the root (Fig. 7A), stem (Fig. 7B), and leaf (Fig. 7C) of WT and *SlWRKY80* transgenic lines showed the same trend. *SlWRKY80*-overexpressing lines had a significantly reduced Na^+/K^+ ratio, promoting K^+ absorption and Na^+ outflow. Through direct comparison of the evolutionary trees of tomato *SINHX* and *Arabidopsis AtNHX*, it was found that the genetic relationship between *AtNHX1*, *AtNHX2*, and *SINHX4* was as high as 82%, with the closest genetic relationship (Supplementary Data Fig. S4C), and the expression level of *SINHX4* significantly increased in response to salt stress [73]. Reports have shown that *AtNHX1* and *AtNHX2* are located in vacuolar membranes and are responsible for regulating the transport of Na^+ and K^+ [8]. At the transcriptional level, we measured the relative expression of *SINHX4*, which regulates ion transport. The *SlWRKY80*-overexpressing lines showed significantly increased relative expression of *SINHX4*, while the *SlWRKY80* knockout lines showed the opposite effect. At the same time, the *SlWRKY80*-overexpressing lines further enhanced the promotion of *SINHX4* expression after saline-alkali stress (Supplementary Data Fig. S10).

To further explore the mechanism of *SlWRKY80* regulating Na^+ and K^+ homeostasis, we conducted a variety of studies on the functional genes regulating Na^+ and K^+ transport, and finally found that *SINHX4* may be directly regulated by *SlWRKY80*. By analyzing the 3000-bp sequence upstream of the *SINHX4* promoter, we found a W-box. Then we confirmed that *SlWRKY80* can directly bind to the W-box in the *SINHX4* promoter by yeast

single hybridization (Y1H) (Fig. 7D), ChIP-qPCR (Fig. 7E), and EMSA (Fig. 7F) *in vivo* and *in vitro*. To sum up, we concluded that under saline-alkali stress the expression of *SlWRKY80* was enhanced, and then regulated the osmotic stress of tomato by regulating *SINHX4* to further promote the absorption of K^+ and the efflux of Na^+ to reduce Na^+/K^+ .

Interaction between *SlWRKY80* and *SlJAZ1* inhibited the regulation of *SlSPDS2* and *SINHX4* by *SlWRKY80*

To further verify the effect of interaction between *SlWRKY80* and *SlJAZ1* on the regulation of downstream functional genes, we proved that *SlWRKY80* positively regulated *SlSPDS2* and *SINHX4* through the dual luciferase test, which was also consistent with the result of ChIP-qPCR (Fig. 6C and Fig. 7E), but when *SlJAZ1* was present the regulation by *SlWRKY80* of downstream *SlSPDS2* and *SINHX4* decreased significantly (Fig. 8A and B). The interaction of *SlJAZ1* and *SlWRKY80* inhibited the regulation by *SlWRKY80* of downstream *SlSPDS2* and *SINHX4*. In addition, the JA synthesis mutant *spr8* is a mutant material based on the CM WT tomato, so in order to further validate this result at the transcriptional level, we treated CM WT tomato as follows: spraying 22.5 $\mu\text{mol/l}$ MeJA or spraying 22.5 $\mu\text{mol/l}$ fluridone, a JA synthesis inhibitor, was used to JA synthesize mutant *spr8*. The use of JA synthesis mutant *spr8* can infer whether *SlWRKY80* affects the entire pathway of JA synthesis or a certain segment of JA synthesis with *spr8* as the node, so it has certain reference significance. The relative expression levels of *SlWRKY80*, *SlSPDS2*, and *SINHX4* in the fluridone group and the *spr8* group were significantly lower than those in the WT group, while the relative expression trend of *SlJAZ1* was opposite (Fig. 8C). Exogenous MeJA promoted the expression of *SlWRKY80*, *SlSPDS2*,




Fig. 5. Overexpression of SIWRKY80 increases endogenous MeJA content and interacts with transcriptional suppressor SIJAZ1. **A** Endogenous MeJA content of WT and SIWRKY80 transgenic lines pre- and 24 h-post-processing. **B** Content of endogenous MeJA in WT, WT + MeJA, 80CR-3, and 80CR-3 + MeJA pre- and 24 h-post-processing. According to the LSD test, significant differences are indicated by lowercase letters ($P < 0.05$). **C** Self-activation verification of SIWRKY80 and verification of the interaction relationship between SIWRKY80 and tomato SIJAZ family through the Y2H test *in vitro*. **D, E** Interaction between SIWRKY80 and SIJAZ1 through the BiFC test and LCI *in vivo*. **F** Verification of the interaction relationship between SIWRKY80 and SIJAZ1 through a pull-down test *in vitro*.

and *SlNHX4*, while inhibiting *SIJAZ1* led to a significant 70.6% decrease in *SIJAZ1* expression. This also confirms the negative regulatory relationship between *SIJAZ1* and *SIWRKY80* at the transcriptional level.

To sum up, we can propose a model in which, under normal conditions, *SIWRKY80* protein interacts with *SIJAZ1* protein, and *SIWRKY80* is bound by *SIJAZ1*. Under 300 mM saline-alkali stress, exogenous spraying of 22.5 $\mu\text{mol/l}$ MeJA can significantly increase the content of endogenous MeJA and JA and accelerate the decomposition of *SIJAZ1*, which weakens or relieves the inhibitory effect of *SIJAZ1* on *SIWRKY80*, thus releasing a large number of *SIWRKY80* proteins to bind to the promoters of *SISPD52* and *SlNHX4* and activate the expression of these two downstream factors, hence promoting the synthesis of Spd and the homeostasis of Na^+ and K^+ , thus regulating saline-alkali stress (Fig. 9).

Discussion

Saline-alkali stress resistance in tomato can be improved by exogenous methyl jasmonate at certain concentrations

JA plays a vital role, such as mechanical damage, disease, insect damage, drought, salt stress, high and low temperature [42, 71].

Arabidopsis thaliana [46], maize (*Zea mays* L.) [71] and *Dioscorea zingiberensis* [51] showed significantly increased internal JA content to enhance salt resistance. By exogenously applying JA to soybean (*Glycine max*), salt stress could be alleviated [54]. In addition, using MeJA exogenously can increase salt tolerance of plants by maintaining ROS or ion homeostasis [69].

In this study, exogenous MeJA increased resistance to saline-alkali-treated tomato at low concentration, and decreased resistance at high concentration. A 300 mM solution of saline-alkali was applied to tomato plants of the same growth potential, and different concentrations of MeJA were sprayed externally. When the concentration of MeJA was 22.5 $\mu\text{mol/l}$, the plant showed significantly higher morphological and physiological indicators (Fig. 1 and Supplementary Data Fig. S1). A dose of exogenous MeJA of 22.5 $\mu\text{mol/l}$ significantly improves tomato protection against saline-alkali stress. When concentrations of exogenous MeJA are low, as a signaling molecule MeJA activates transcription factors in plants under stress [31]. In contrast, when exogenous MeJA is applied at high levels the plant absorbs excess MeJA, increasing its osmotic potential and causing osmotic stress. In addition, excessive exogenous MeJA can affect the balance between endogenous hormones, leading to abnormal regulation of plant hormones.




Fig. 6. Binding of *SlWRKY80* to the promoter region of *SlSPDS2* and promotion of *Spd* synthesis. **A** *Spd* content in *SlWRKY80* transgenic tomato materials before and after saline–alkali treatment. **B** Y1H assay. **C** *SlWRKY80* interaction with the *SlSPDS2* promoter confirmed by ChIP–qPCR assay. The three W-boxes within the 3000-bp upstream promoter of *SlSPDS2* are denoted as pro-*SlSPDS2*-P1, pro-*SlSPDS2*-P2, and pro-*SlSPDS2*-P3. According to the LSD test, significant differences are indicated by lowercase letters ($P < 0.05$), and values indicate the mean across three biological replicates. **D** *SlWRKY80* interacts with the *SlSPDS2* promoter *in vitro*, as shown by EMSA.

SlWRKY80 senses methyl jasmonate and saline–alkali signals and actively regulates saline–alkali stress

An important role of WRKY genes is to regulate the response of plants to biotic and abiotic stresses. The group III subfamily of tomato *SlWRKY* mainly responds to abiotic stresses in tomato [12, 24]. Toward a deeper understanding of how MeJA protects tomato against saline–alkali stress, we found *SlWRKY80*, located in group III of *SlWRKY* through transcriptome analysis before and after saline–alkali stress (Fig. 2A). *SlWRKY80* expression was highest in the S+M group, followed by the

S group, and both were significantly higher than in the WT group (Fig. 2B). Meanwhile, the GUS staining results of pCAMBIA1391-pro-*SlWRKY80* transgenic material also showed that the promoter of *SlWRKY80* responded to MeJA and saline–alkali stress (Fig. 2C). Similarly, SA induced the mulberry *MiWRKY53* promoter to activate the resistance of *MiWRKY53* to *Pseudomonas syringae* [43].

To better explore the mechanism of *SlWRKY80* under saline–alkali stress, we obtained *SlWRKY80*-overexpressing lines (Supplementary Data Fig. S5) and knockout lines (Supplementary Data Fig. S6) by transgenic methods. After 300 mM




Fig. 7. By binding to the promoter of *SINHX4*, *SIWRKY80*-overexpressing lines decrease the Na⁺/K⁺ ratio in tomato. **A-C** Na⁺ and K⁺ contents and Na⁺/K⁺ ratio of root, stem, and leaf. According to the LSD test, significant differences are indicated by lowercase letters ($P < 0.05$), and values indicate the mean across three biological replicates. **D** Y1H assay. **E** ChIP-qPCR. **F** *SIWRKY80* interacts with the *SINHX4* promoter *in vitro*, as shown by EMSA.

saline-alkali treatment, the growth of *SIWRKY80*-overexpressing lines was the best, and the growth of *SIWRKY80* knockout lines was the worst (Fig. 3A). DNB staining (Fig. 3C) and NBT staining (Fig. 3D) also showed that the staining of *SIWRKY80*-overexpressing lines was the lightest, and the staining of *SIWRKY80* knockout lines was the deepest, indicating that the *SIWRKY80*-overexpressing lines removed more ROS, which was also consistent with the results of SOD activity, POD activity, CAT activity (Fig. 3G), and other indicators on the eighth day of treatment; the other morphological and physiological indexes were the same. Above all, *SIWRKY80* can actively regulate saline-alkali stress and the *SIWRKY80* promoter was responsive to exogenous MeJA (22.5 $\mu\text{mol/l}$), which stimulated

the expression of *SIWRKY80* and further regulated saline-alkali stress.

Another interesting phenomenon is that exogenous MeJA (22.5 $\mu\text{mol/l}$) spraying of *SIWRKY80* knockout lines (80CR-3) can significantly increase the seedling strength index, SOD activity, POD activity, and CAT activity (Fig. 4), indicating that exogenous MeJA (22.5 $\mu\text{mol/l}$) spraying can significantly reduce the sensitivity of *SIWRKY80* knockout lines to saline-alkali stress. This phenomenon may be due to the activation of multiple metabolic pathways by exogenous MeJA as a signal molecule; *SIWRKY80* is only an important transcription factor in exogenous MeJA resistance to saline-alkali stress. Similarly, The CmGST family of genes, including *CaGSTU3*, *CaGSTU7*, and others is




Fig. 8. Results of the dual luciferase reporter assay. **A** A control plasmid, 35S:REN, was co-infiltrated as an interfering plasmid into *N. benthamiana* leaves. **B** Interaction between SIWRKY80 and SIJAZ1, resulting in the inhibition of SIWRKY80 regulation of SISPDS2 (-P1/P2/P3) and SINHX4, respectively. LUC/REN ratios were used to determine the ability of SIWRKY80 and SIJAZ1 to activate the reporter LUC gene. **C** Relative expression levels of SIWRKY80, SISPDS2, SINHX4, and SIJAZ1 after exogenous spraying of MeJA or fluridone; *spr8* mutants were measured against the background of WT tomato CM. According to the LSD test, significant differences are indicated by lowercase letters ($P < 0.05$), and values indicate the mean across three biological replicates.

believed to contribute to pumpkins' cold resistance [1]. Therefore, a new reference for studying JA stress resistance mechanisms is provided by this study.

SIWRKY80 directly combines with SISPDS2 and SINHX4 promoters to regulate spermidine synthesis and Na^+/K^+ homeostasis

Spd is a polyamine compound that helps plants adapt to abiotic stress and grow and develop [34, 64]. Exogenous spraying of Spd protects membrane lipids from peroxidation, regulates polyamine metabolism, and strengthens the antioxidant system in tomato [20, 33, 67]. Elevated endogenous Spd levels in tomato plants have been shown to significantly enhance their tolerance to saline-alkali stress [56]. In addition, saline-alkali stress primarily disrupts cellular ion homeostasis, emphasizing the need to understand the mechanisms of Na^+ absorption and transport in plants and identify candidate genes that promote

ion homeostasis to enhance crop salt tolerance [9]. Therefore, we wondered whether SIWRKY80 actively regulated saline-alkali stress was related to Spd synthesis and Na^+/K^+ homeostasis or not.

We discovered that SIWRKY80-overexpressing lines had significantly increased the Spd content, which was further enhanced after 8 days of saline-alkali treatment (Fig. 5A). Similar results were found in different tissue parts. The contents of K^+ in root (Fig. 6A), stem (Fig. 6B), leaf (Fig. 6C) and other tissue parts of SIWRKY80-overexpressing lines were significantly higher than those of WT. Likewise, the opposite was true for SIWRKY80 knockout lines. We found the Spd synthesis gene (SISPDS2) and the SINHX4 gene related to Na^+/K^+ transport through the transcriptome and from the literature, with confirmation through *in vivo* and *in vitro* experiments (Y1H, EMSA, ChIP-qPCR) that SIWRKY80 binds to SISPDS2 (Fig. 5B-D) and SINHX4 promoters (Fig. 6D-F), respectively. In our previous research, we found that




Fig. 9. A working model for the saline-alkali reaction mediated by *SIWRKY80* through the JA pathway and the regulation of *SISPDS2* and *SINHX4* in tomato. The CGTCA motif is a MeJA-binding element.

overexpression of *SISPDS2* [56] and *SISPMS* [63] can increase the endogenous free polyamine content in tomato seeds. This leads to the regulation of ion balance, the antioxidant enzyme system, and osmotic regulators under saline-alkali stress, thus enhancing the resistance of tomato seeds to saline-alkali stress.

Following saline-alkali treatment, the Na^+/K^+ ratio in the *SIWRKY80*-overexpressing lines was notably lower compared with the WT, whereas the opposite was observed in the *SIWRKY80* knockout lines (Fig. 7A–C). Additionally, *SIWRKY80* exhibited active regulation of saline-alkali stress (Fig. 4). Consequently, the *SIWRKY80*-overexpressing lines may actively respond to saline-alkali stress through its involvement in the transportation of Na^+ and K^+ . In addition, the genetic relationship between *AtNHX1*, *AtNHX2*, and *SINHX4* was as high as 82% (Supplementary Data Fig. S4C), the genetic relationship between *AtNHX1*, *AtNHX2*, and *SINHX4* is very close, so *SINHX4* may have similar physiological functions as *AtNHX1* and *AtNHX2*, and the expression level of *SINHX4* significantly increased in response to salt stress [73]. *AtNHX1* and *AtNHX2* are located in vacuolar membranes and are responsible for regulating the transport of Na^+ and K^+ [8]. The expression level of *SINHX4* was significantly upregulated in the saline-alkali treatment and in *SIWRKY80*-overexpressing lines in this experiment. Finally, this experiment validated the upstream relationship between *SIWRKY80* and *SINHX4* through molecular experiments such as Y1H (Fig. 7D) and EMSA (Fig. 7F). Under salt stress, *NHX* promoted the entry of Na^+ into the vacuole and the absorption of K^+ in tomato [4, 8]. Therefore, an increase in the endogenous Spd content and decrease in the Na^+/K^+ ratio are helpful to improve the saline-alkali resistance of tomato. Thus, we conclude that *SIWRKY80* promotes the synthesis of Spd and Na^+/K^+ homeostasis by interacting with downstream *SISPDS2* and *SINHX4* to actively regulate saline-alkali stress.

SIJAZ1 is inhibited in saline-alkali stress and releases more *SIWRKY80* when interacting with it

Studies have shown that MeJA and JA influence plant growth, development, and stress responses. Increased endogenous JA can reduce salt damage in wheat [47], rapeseed [28], rice [6], and other crops by activating genes involved in the JA signaling pathway.

Salt damage can be alleviated by JA in wheat [47] and rapeseed [28]. Endogenous JA content increased and its signal transduction was activated when salt damage occurred [69]. In our study, after 24 h of saline-alkali stress, the endogenous MeJA and JA contents of tomato were significantly increased. *SIWRKY80*-overexpressing lines had significantly higher MeJA and JA contents than WT, whereas those of *SIWRKY80* knockout lines were significantly lower (Fig. 7A), indicating that the *SIWRKY80*-overexpressing lines can show significantly increased endogenous MeJA and JA contents in tomato in saline-alkali stress, while the knockout lines exhibited contrary results. Under saline-alkali stress, exogenous application of 22.5 $\mu\text{mol/l}$ MeJA to the 80CR-3 knockout lines significantly increased the endogenous MeJA and JA contents of tomato seedlings after 24 h (Fig. 7B), and the relative expression levels of JA synthesis-related genes such as *SILOxD* and *SILOc* also significantly increased (Supplementary Data Fig. S11), indicating that exogenous MeJA can significantly increase the endogenous MeJA and JA contents of tomato seedlings. In the results of 80CR-3 and 80CR-3 + MeJA under saline-alkali stress (Fig. 4), GUS staining results (Fig. 2C), Spd content (Fig. 5A), and Na^+/K^+ transport (Fig. 6A–C), both saline-alkali stress and exogenous MeJA treatment showed that they activated the promoter of *SIWRKY80* and led to its expression. *SIWRKY80* overexpression also promoted the expression of *SISPDS2* (Supplementary Data Fig. S9) and *SINHX4* (Supplementary Data Fig. S10). We also affirmed that *SIWRKY80* can directly regulate the promoters of *SISPDS2* and *SINHX4*, which implies that saline-alkali stress and MeJA exogenous spray actively regulate saline-alkali stress by increasing the endogenous MeJA and JA levels.

Allele oxidase synthase (AOS), allele oxycyclinase (AOC), and 12-oxo plant dienoic acid reductase (OPR) are the rate-limiting enzymes in the JA biosynthetic pathway, while coronatine insensitive1 (COI1) and JAZ proteins are two important receptors in the JA biosynthetic pathway [59]. JAZ protein contains the jasmonate Zim domain. As an inhibitor, the JA signaling pathway can be connected to other signaling pathways through combinations with transcription factors or other coenzyme proteins [5]. There are 13 genes in the tomato *SIJAZ1* family, so we verified the interaction of *SIWRKY80* with *SIJAZ1*, *SIJAZ2*, and *SIJAZ5* at protein level through the Y2H assay (Fig. 7C). We verified it by LCI (Fig. 7D), BiFC (Fig. 7E), and pull-down assays (Fig. 7F) *in vivo* and *in vitro*. We found that

only *SlJAZ1* could get positive results under verification by these molecular means. Therefore, we conclude that there is a protein-level interaction between *SlWRKY80* and *SlJAZ1*.

An SCF^{CO1} complex is formed when CO1 combines with JAZ protein, which results in JAZ protein degradation by the 26S proteasome, then releases transcription factors that interact with JAZ, thereby activating the expression of JA-responsive genes. Therefore, JA content is negatively correlated with the expression of *SlJAZs* [30, 50]. The same results were obtained in our study. When the synthesis of JA in tomato was inhibited, the expression of *SlJAZ1* was significantly upregulated, while the relative expression of *SlWRKY80*, *SlSPDS2*, and *SlNHX4* was opposite (Fig. 8C), indicating that the increase of endogenous JA content inhibited the expression of *SlJAZ1*. The interaction between *SlJAZ1* as a transcriptional suppressor gene and *SlWRKY80* will also be weakened, thus releasing more *SlWRKY80* protein. This result was also proved by the dual luciferase test (Fig. 8B).

To sum up, saline-alkali stress or exogenous spraying with a certain concentration of MeJA could increase the contents of endogenous MeJA and JA in tomato. As a signal molecule, MeJA combined with the *SlWRKY80* promoter to promote the expression of *SlWRKY80*. On the other hand, the increase of endogenous JA content inhibited the expression of *SlJAZ1* and further released *SlWRKY80* protein, interacting with *SlJAZ1*. At the same time, *SlWRKY80* combined with the promoters of *SlSPDS2* and *SlNHX4* to promote Spd synthesis and Na⁺/K⁺ homeostasis, thus actively regulating saline-alkali stress.

Materials and methods

Plant materials

To verify the optimal concentration of exogenous MeJA under 300 mM saline-alkali stress, WT 'Ailsa Craig' (AC) tomatoes were used, and AC was also used as the background material to construct *SlWRKY80*-overexpressing lines (80OE-1 and 80OE-3) and *SlWRKY80* knockout lines (80CR-3 and 80CR-4). In addition, the JA synthesis mutant *spr8* (Solyc03g122340) used in this experiment was provided by Professor Li Chuanyou (Institute of Genetics and Developmental Biology, Chinese Academy of Sciences), and the background material for the knockout lines was CM WT tomato.

Obtaining and identification of *SlWRKY80*-overexpressing and knockout lines

Overexpression vectors were constructed from tomato AC and pHellsgate2 (CaMV35S promoter driver) overexpressing *SlWRKY80* by cloning the CDS with SmaI and KpnI (an HA label was added to the pHellsgate2 vector, which was modified by Zhan Xiangqiang, School of Horticulture, Northwest A&F University). *SlWRKY80* target information was predicted through CRISPR RGEN Tools (<http://www.rgenome.net/casdesigner/result?hash=49d4ef08e39c96a14781bc8f463be7f6>) and constructing CRISPR/Cas9:*SlWRKY80*. Sequencing was performed by Sangon Biotech (Shanghai, China). *SlWRKY80* from different genotypes was compared with DNAMAN v.6 (Lynnon Biosoft, CA, USA). The primers used are listed in Supplementary Data Table S1.

The above-mentioned recombinant vectors were utilized to obtain transgenic materials via *Agrobacterium*-mediated infection [65]. Identification was performed using specific primers (Supplementary Data Table S1), and positive materials were retained for seed collection and continued propagation. The T₂ generations of homozygous overexpressing (80OE-1, 80OE-3) and knockout lines (80CR-3, 80CR-4) were obtained by self-crossing to obtained transgenic lines.

Plant growth conditions

This experiment was conducted in a growth chamber (model GXZ-5000E, China Ningbo Southeast Instrument Co., Ltd) under controlled conditions, with a daily light period of 12 h at 200 μmol m⁻² s⁻¹, temperature 25°C (day) and 20°C (night), humidity set to 60%. At the age of 15 days, we selected seedlings with consistent growth and planted them in a nutrient bowl and treated them at the age of 35 days (approximately five true leaves).

Treatments

In the saline-alkali tolerance assay, a 300 mM composite saline-alkali solution (NaCl:Na₂SO₄:NaHCO₃:Na₂CO₃ = 1:9:9:1, molar content ratio, pH = 8.6 ± 0.1) was used based on extensive saline-alkali stress resistance testing conducted by our research team [33, 56, 67, 68].

We verified the effect of exogenous MeJA at different concentrations on tomato stress resistance under saline-alkali stress. In the present study we sprayed MeJA at 0, 12.5, 22.5, 45, and 90 μmol/l in WT as a preliminary experiment and based on the report of Min *et al.* [42]. AC tomato seedlings at 35 days of age were irrigated with 100 mL of 300 mM saline-alkali solution, while external MeJA was applied by spraying. The concentrations of external MeJA were 0 (WT), 12.5, 22.5, 45, and 90 μmol/l. Each sprayed plant was 5 cm away from the tomato once, front, back, left, right. On average, ~5 ml of exogenous MeJA was sprayed per plant.

To investigate the effects of saline-alkali stress and exogenous MeJA on *SlWRKY80* expression and the promoter of *SlWRKY80*, we set up three groups of treatments, namely WT (control), S, and S + M. The WT (control) plants were irrigated with distilled water (100 ml) and the S (saline-alkali treatment) plants were irrigated with 300 mM saline-alkali (100 ml), while the S + M (saline-alkali and MeJA co-treatment) plants were irrigated with 300 mM saline-alkali solution (100 ml) and sprayed with 22.5 μmol/l of exogenous MeJA (CAS No. 39924-52-2, Sigma-Aldrich, USA). AC seedlings were treated at 0, 3, 6, 12, and 24 h, and the third leaf from the top of the tomato plant to measure the relative expression level of *SlWRKY80*. The pCAMBIA1391-pro-*SlWRKY80* material was subjected to GUS staining at 24 h of treatment.

A saline-alkali stress study was conducted to verify the function of *SlWRKY80*. We subjected AC (WT), *SlWRKY80*-overexpressing, and knockout lines from the same growth period to 300 mM saline-alkali treatment, and observed the phenotype 8 days later. We verified whether exogenous application of MeJA reduced the sensitivity of the 80CR-3 lines to saline-alkali stress. We selected AC and 80CR-3 tomato seedlings (35 days age) and grouped them into three treatments, namely WT, 80CR-3, and 80CR-3 + MeJA. Using AC tomato as WT, we poured 100 ml of saline-alkali solution on all three groups, and sprayed water (80CR-3) and an equal volume of 22.5 μmol/l MeJA (80CR-3 + MeJA) on 80CR-3. This experiment lasted for 8 days, during which the differences between each group were observed on a daily basis.

Further verification of the relationship between *SlWRKY80*, *SlSPDS2*, *SlNHX4*, and *SlJAZ1* was performed. We were fortunate enough to obtain a mutant *spr8* (*SlLoxD* gene mutant tomato material) synthesized from JA (using CM WT tomato as the background material). Based on this, we used CM WT tomato as the material, treated it with 300 mM saline-alkali, and applied 22.5 μmol/l MeJA and 22.5 μmol/l fluridone (JA synthesis inhibitor) (CAS No. CF5275, Beijing, China), externally, while spraying the same volume of ddH₂O on *spr8*.

RNA extraction and real-time quantitative PCR

Using 35-day-old WT, 80OE-1, 80OE-3, 80CR-3, and 80CR-4 tomato seedlings, we selected the third leaf from the top of the tomato plant, extracted RNA from the stem tissue and the entire root system, and measured the relative expression level of *SlWRKY80*. RNA extraction and first-strand cDNA synthesis and RT-qPCR were carried out following the protocol established by Xu et al. [65] and Livak et al. [37]. Related primers are provided in [Supplementary Data Table S2](#).

In the experiments using saline-alkali treatment (S) and saline-alkali and MeJA co-treatment (S+M), we collected leaf tissue samples at 0, 6, and 12 h of treatment, and extracted RNA to measure *SlWRKY80* expression. The sampling position was the third blade from the top.

Determination of morphological and physiological indicators

The phenotypes of the WT and *SlWRKY80* transgenic lines treated with a 300 mM saline-alkali solution were observed, and various physiological parameters were determined on the eighth day. A ruler, vernier caliper, and root scanner (Perfection V700N, Epson Co., Ltd, China) were used to measure aboveground morphological indicators [68]. The formula for calculating the sound seedling index was: (stem diameter/plant height + root dry weight/aboveground dry weight) × whole plant dry weight.

The physiological indicators included the activities of SOD, POD, and CAT, proline content, malondialdehyde content, and so on. A reagent kit was used to measure the physiological indicators (Nanjing Jiancheng Biotechnology Research, Nanjing, China).

Histochemical GUS activity assay

To clarify the response of the promoter of *SlWRKY80* to MeJA, JA-Ile and saline-alkali, the pCAMBIA1391-pro-*SlWRKY80* vector was constructed using HindIII and SalI as restriction endonuclease sites and transferred into AC tomato plants through genetic transformation, and pCAMBIA1391-pro-*SlWRKY80*-positive material was obtained. The relevant primers are shown in [Supplementary Data Table S1](#).

Untreated pCAMBIA1391-pro-*SlWRKY80*-positive material was used as a control in this experiment. We performed S, M, S+M processing separately. After 24 h of treatment, GUS staining was performed referring to Liang et al. [29] (The decolorization process after GUS staining ends when the AC leaves are decolorized until colorless).

Subcellular localization of *SlWRKY80*

The CDS of *SlWRKY80*, excluding the stop codon, was cloned into the pBI121-GFP vector at the SacI and BamHI restriction sites, resulting in the generation of the pBI121-*SlWRKY80*-GFP construct. Additionally, the pBI121-GFP construct was produced using the same SacI and BamHI restriction sites. The primers used are mentioned in [Supplementary Data Table S1](#). For subcellular localization, the pCMV-C-mCherry vector (Beyotime, Shanghai, China) was employed. The plasmids were introduced into *Agrobacterium tumefaciens* GV3101 utilizing the freeze-thaw method. Transient transformation of *Nicotiana benthamiana* leaf epidermal cells was subsequently conducted. Detailed operation methods can be found in Xu et al. [66].

Determination of free spermidine content

On the eighth day of treatment, the second leaf from the top was sampled, with three biological replicates from each group.

The determination of free Spd content was carried out using the method of Wang et al. [56].

Determination of Na⁺ and K⁺ contents

On the eighth day of treatment, samples were collected from roots, stems, and leaves. The entire root system of the tomato was selected, and the stem was selected at a distance of 1–3 cm from the root system. The second true leaf from the top was selected, and each group of treatments underwent three biological replicates. Na⁺ and K⁺ contents were determined using Wang et al.'s method [58].

Determination of endogenous hormone content

JA and MeJA contents in tomato leaves were determined by taking samples from the second true leaf from the top 0 and 24 h post-treatment. The determination of endogenous hormone content was carried out using the method of Xu et al. [66].

Yeast one-hybrid assay

SlWRKY80 cDNA was sequenced and cloned into pGADT7 vector for construction of prey-*SlWRKY80*. Three distinct fragments of the *SlSPDS2* promoter, along with the *SINH4* promoter, were individually inserted into the pAbAi vector to construct pBait-reporter vectors. Upon completing vector construction, the Y1H procedure was conducted in accordance with the methodology outlined by Liang et al. [29]. [Supplementary Data Table S1](#) lists the related primers.

Chromatin immunoprecipitation qPCR assay

Total protein extracts from WT and 80OE-13 leaves on pre- and post-saline-alkali plants were subjected separately to ChIP assays using a ChIP kit (Beyotime). HA antibodies were used to immunoprecipitate DNA-protein complexes (Sigma). After the precipitated complexes were recovered, ChIP-qPCR assays using the primers provided in [Supplementary Data Table S2](#) were performed.

Electrophoretic mobility shift assay

The CDS of *SlWRKY80* was cloned into the pGEX4T-1 expression vector using the EcoRI and SalI sites to produce a glutathione S-transferase fusion protein. Related primers can be found in [Supplementary Data Table S1](#). The fusion protein was then transformed into *Escherichia coli* line BL21, and induction was carried out using constant shaking at 200 revolutions/min and a temperature of 28°C for 8 h with an IPTG concentration of 0.5 mM. A method based on glutathione Sepharose beads (635608, Takara) was then used to purify the fusion protein. In vitro synthesized the biotin-labeled *SlSPDS2* and *SINH4* promoter oligonucleotide probes, which are listed in [Supplementary Data Table S1](#). EMSAs were performed utilizing a LightShift Chemiluminescent EMSA Kit (Beyotime). The mutation and cold probes employed for EMSA can be found in [Supplementary Data Table S3](#).

Yeast two-hybrid assay

To assess the self-activation of *SlWRKY80*, we divided it into two fragments consisting of a conserved domain, which were subsequently cloned separately into the yeast pGBKT7 vector, with BD-*SlWRKY80*_{202-819bp} and BD-*SlWRKY80*_{1-819bp} cleavage sites. [Supplementary Data Table S1](#) lists the primer sequences. BD-*SlWRKY80*_{202-819bp}, BD-*SlWRKY80*_{1-819bp}, and pGADT7 were transformed into Y2H. After transformation, 50 μl of the transformed cells was placed onto solid-state two-deficient DDO (SD/-Trp/-Leu) and four-deficient QDO (SD/-Trp/-Leu/-His/

–Ade) media. The media were incubated for 3–5 days at 28°C, and yeast growth was recorded. This showed that BD-SIWRKY80_{202-819bp} was a non-self-activating fragment containing the structural domain of SIWRKY80.

The CDSs of 13 genes from the SIJAZ family in tomato were cloned into yeast pGADT7 vector with NdeI and XhoI cleavage sites, and then co-transformed with BD-SIWRKY80_{202-819bp} into Y2H. After transformation, 50 µl of the transformed cells was placed onto solid-state media of two-deficient DDO and four-deficient QDO. The media were placed in an incubator at 28°C for 3–5 days.

Bimolecular fluorescence complementation assay

The full-length CDSs of SIWRKY80 and SIJAZ1 (excluding the termination codon) were constructed in the pSPYNE and pSPYCE vectors, respectively. [Supplementary Data Table S1](#) provides the primers. Subsequently, the constructed vector was sequenced and validated, and the plasmid was transferred into *Agrobacterium* GV3101. The empty plasmids pSPYNE and pSPYCE, along with the recombinant plasmid-containing *Agrobacterium*, were diluted with MES (OD₆₀₀ = 1.0). Then, *Agrobacterium* containing pSPYNE-X (Vec or SIWRKY80) and pSPYCE (Vec or SIJAZ1) was mixed in a 1:1 volume ratio and left to stand in a dark environment at 28°C for 2 h before injection into tobacco leaves. After 48 h, the infected tissue was examined using a confocal microscope (LAS X, Leica, Mannheim, Germany). The images were then subjected to postprocessing using Leica LAS X software (v.3.7.2). The scale used in this experiment was 20 µm.

Luciferase complementary imaging experiment

The CDS of SIWRKY80 was constructed into the N-terminus of JW771 (N-Luc), and SIJAZ1 was constructed into the C-terminus of JW772 (C-Luc). [Supplementary Data Table S1](#) provides the primers. Subsequently, the recombinant plasmid with accurate sequencing was transferred into GV3101. The empty plasmids JW771 and JW772, as well as the recombinant plasmid-containing *Agrobacterium*, were diluted with infection buffer (OD₆₀₀ = 1.0). Then, *Agrobacterium* containing JW771-X (Vec or SIWRKY80) and JW772 (Vec or SIJAZ1) was mixed in a 1:1 volume ratio and left to stand in a dark environment at 28°C for 2 h before injection into tobacco leaves. After 48 h, 0.5 mM D-luciferin was evenly applied to the back of tobacco leaves, and photographs were taken using a plant living molecule labeling imaging system (CCD). The scale was 20 µm.

Pull-down assay

Induction and purification of the GST–SIWRKY80 fusion protein is described above in the section ‘Electrophoretic mobility shift assay’. Cloning of the CDS of SIJAZ1 into pMAL-C2x was performed using EcoRI and HindIII. The expression vector was generated using relevant primers ([Supplementary Data Table S1](#)) to produce the MBP-SIJAZ1 fusion protein. Next, the *E. coli* BL21 line was transformed with the fusion protein and grown by shaking the culture at 200 revolutions/min and 28°C for 8 h, with induction using IPTG at a concentration of 0.5 mM. Subsequent removal of the medium was performed by centrifugation (temperature 4°C, speed 4000 rpm, duration 10 min), and the supernatant was discarded. The remaining substance was resuspended in 1 × PBS (pH = 8.0) and subsequently subjected to ultrasound treatment on ice for 15 min. The solution was then centrifuged again (temperature 4°C, speed 4000 rpm, duration 10 min), and the supernatant was collected. A maltose binding protein label protein purification

kit was used to further purify the obtained MBP-SIJAZ1 fusion protein (Abbkine Biotechnology Co., Ltd, Wuhan, China).

The GST pull-down analysis method used was as described by Zhang et al. [72].

Dual-luciferase analysis

The full-length CDS of SIWRKY80, excluding the termination codon, was cloned into the pGreenII-002962-SK vector using SacI and KpnI cleavage sites. The *SISPDS2* promoter has three promoter segments while the *SINH4* promoter has one promoter segment containing the W-box. They were constructed in the pGreen-II-0800 vector, with KpnI and NcoI enzyme cleavage sites. The primers used are listed in [Supplementary Data Table S1](#). The dual luciferase test operation was based on the method of Liang et al. [29]. The tobacco plants were then incubated for 72 h before fluorescence detection was performed using the Dual Lucifera Reporter Assay System (E1910, Promega, USA).

Statistical analysis

The LSD test, based on DPS7.5, was used to detect significant differences between three replicates of each experiment.

Acknowledgements

Several funding sources were used to complete this work, including Shaanxi Province’s Scientific & Technological Innovative Research Team (2021TD-34) and the China Agriculture Research System (CARS-23-D06). We thank Professor Chuanyou Li and Lei Deng for providing *spr8* and Professor Xiangqiang Zhan for providing the overexpression vector. We thank Jing Zhang for assistance in the MeJA and JA analysis, and Yangyang Yuan for helping in the use of a confocal laser scanning microscope. Anti-Flag was purchased from Sangon Biotech; anti-GST and anti-MBP antibodies were purchased from Abcam.

Author contributions

X.H., G.L., C.S., and X.L. conceived and designed the experiments. C.S., G.L., and X.H. wrote the paper. C.S. and X.L. performed the experiments. C.S., X.L., G.C., H.Z., and G.L. provided advice related to the research. Author A.K. provided great assistance in polishing the manuscript. All authors read and approved the manuscript for submission.

Data availability

The authors affirm that all the information required to substantiate the findings of the study is provided in both the paper and the supplementary materials, or can be acquired by contacting the corresponding author.

Conflict of interest

The authors declare no competing commercial or financial interest.

Supplementary data

[Supplementary data](#) is available at *Horticulture Research* online.

Accession numbers

A database of online sequence data was used to obtain sequence data (https://solgenomics.net/search/locus?tdsourcetag=spcqq_aimsg).

The gene accession numbers used in this study can be found below: Solyc03g095770.2 (SlWRKY80), Solyc01g098190.3 (SlNHX4), (SlSPDS2), Solyc07g042170.3 (SlJAZ1), Solyc12g009220.2 (SlJAZ2), Solyc03g122190.3 (SlJAZ3), Solyc12g049400.2 (SlJAZ4), Solyc03g118540.3 (SlJAZ5), Solyc01g005440.3 (SlJAZ6), Solyc11g011030.2 (SlJAZ7), Solyc06g068930.2 (SlJAZ8), Solyc08g036640.3 (SlJAZ9), Solyc08g036620.3 (SlJAZ10), Solyc08g036660.3 (SlJAZ11), Solyc01g009740.2 (SlJAZ12), Solyc01g103600.3 (SlJAZ13), Solyc03g122340.2 (SlLoxD), Solyc02g085730.2 (SlAOC).

References

- Abdul Kayum M, Nath UK, Park JI. et al. Genome-wide identification, characterization, and expression profiling of glutathione S-transferase (GST) family in pumpkin reveals likely role in cold-stress tolerance. *Genes*. 2018;**9**:84
- Abouelsaad I, Renault S. Enhanced oxidative stress in the jasmonic acid-deficient tomato mutant *def-1* exposed to NaCl stress. *J Plant Physiol*. 2018;**226**:136–44
- Ahmad RM, Cheng C, Sheng J. et al. Interruption of jasmonic acid biosynthesis causes differential responses in the roots and shoots of maize seedlings against salt stress. *Int J Mol Sci*. 2019;**20**:6202
- Ali AAM, Romdhane WB, Tarroum M. et al. Analysis of salinity tolerance in tomato introgression lines based on morpho-physiological and molecular traits. *Plants (Basel)*. 2021;**10**:2594
- Andrea C, Isabel M, Gemma FB. et al. A small molecule antagonizes jasmonic acid perception and auxin responses in vascular and nonvascular plants. *Plant Physiol*. 2021;**187**:1399–413
- Asfaw KG, Liu Q, Eghbalian R. et al. The jasmonate biosynthesis gene OsOPR7 can mitigate salinity induced mitochondrial oxidative stress. *Plant Sci*. 2022;**316**:111156
- Bai Y, Sunarti S, Kissoudis C. et al. The role of tomato WRKY genes in plant responses to combined abiotic and biotic stresses. *Front. Plant Sci*. 2018;**9**:801
- Bassil E, Zhang SQ, Gong HJ. et al. Cation specificity of vacuolar NHX-type cation/H⁺ antiporters. *Plant Physiol*. 2019;**179**:616–29
- Basu S, Kumar A, Benazir I. et al. Reassessing the role of ion homeostasis for improving salinity tolerance in crop plants. *Physiol Plant*. 2020;**171**:502–19
- Bo C, Cai RH, Fang X. et al. Transcription factor ZmWRKY20 interacts with ZmWRKY115 to repress expression of *ZmbZIP111* for salt tolerance in maize. *Plant J*. 2022;**111**:1660–75
- Chele KH, Tinte MM, Piater LA. et al. Soil salinity, a serious environmental issue and plant responses: a metabolomics perspective. *Metabolites*. 2021;**11**:724
- Dang F, Lin J, Li Y. et al. SlWRKY30 and SlWRKY81 synergistically modulate tomato immunity to *Ralstonia solanacearum* by directly regulating SlPR-STH2. *Hortic Res*. 2023;**10**:uhad050
- Devkar V, Thirumalaikumar VP, Xue GP. et al. Multifaceted regulatory function of tomato SlTAF1 in the response to salinity stress. *New Phytol*. 2020;**225**:1681–98
- Ding ZJ, Yan JY, Li CX. et al. Transcription factor WRKY46 modulates the development of *Arabidopsis* lateral roots in osmotic/salt stress conditions via regulation of ABA signaling and auxin homeostasis. *Plant J*. 2015;**84**:56–69
- Gao Y, Liu J, Yang F. et al. The WRKY transcription factor WRKY8 promotes resistance to pathogen infection and mediates drought and salt stress tolerance in *Solanum lycopersicum*. *Physiol Plant*. 2019;**168**:98–117
- Garcia-Abellan JO, Fernandez-Garcia N, Lopez-Berenguer C. et al. The tomato *res* mutant which accumulates JA in roots in non-stressed conditions restores cell structure alterations under salinity. *Physiol Plant*. 2015;**155**:296–314
- Hazman M, Hause B, Eiche E. et al. Increased tolerance to salt stress in OPDA-deficient rice ALLENE OXIDE CYCLASE mutants is linked to an increased ROS-scavenging activity. *J Exp Bot*. 2015;**66**:3339–52
- Huertas R, Olías R, Eljakaoui Z. et al. Overexpression of SlSOS2 (SlCIPK24) confers salt tolerance to transgenic tomato. *Plant Cell Environ*. 2012;**35**:1467–82
- He YQ, Zhang XC, Tan YX. et al. Virus-induced gene silencing of SlWRKY79 attenuates salt tolerance in tomato plants. *Agronomy*. 2021;**11**:1519
- Hu XH, Zhang Y, Shi Y. et al. Effect of exogenous spermidine on polyamine content and metabolism in tomato exposed to salinity-alkalinity mixed stress. *Plant Physiol Biochem*. 2012;**57**:200–9
- Hu YR, Chen LG, Wang HP. et al. *Arabidopsis* transcription factor WRKY8 functions antagonistically with its interacting partner VQ9 to modulate salinity stress tolerance. *Plant J*. 2013;**74**:730–45
- Huang SX, Gao YF, Liu JK. et al. Genome-wide analysis of WRKY transcription factors in *Solanum lycopersicum*. *Mol Gen Genomics*. 2012;**287**:495–513
- Jiang Y, Yu D. WRKY57 regulates JAZ genes transcriptionally to compromise 2 *Botrytis cinerea* resistance in *Arabidopsis thaliana*. *Plant Physiol*. 2016;**171**:2771–82
- Jiang JJ, Ma SH, Ye N. et al. WRKY transcription factors in plant responses to stresses. *J Integr Plant Biol*. 2017;**59**:86–101
- Jiang YQ, Deyholos MK. Functional characterization of *Arabidopsis* NaCl-inducible WRKY25 and WRKY33 transcription factors in abiotic stresses. *Plant Mol Biol*. 2009;**69**:91–105
- Jiang YZ, Tong SF, Chen NN. et al. The PalWRKY77 transcription factor negatively regulates salt tolerance and abscisic acid signaling in *Populus*. *Plant J*. 2021;**105**:1258–73
- Kasukabe Y, He L, Nada K. et al. Overexpression of spermidine synthase enhances tolerance to multiple environmental stresses and up-regulates the expression of various stress-regulated genes in transgenic *Arabidopsis thaliana*. *Plant Cell Physiol*. 2004;**45**:712–22
- Kuo HY, Kang FC, Wang YY. Glucosinolate transporter1 involves in salt-induced jasmonate signaling and alleviates the repression of lateral root growth by salt in *Arabidopsis*. *Plant Sci*. 2020;**297**:110487
- Liang Y, Bai JY, Xie ZL. et al. Tomato sucrose transporter SlSUT4 participates in flowering regulation by modulating gibberellin biosynthesis. *Plant Physiol*. 2023;**192**:1080–98
- Li CY, Liu GH, Xu CC. et al. The tomato suppressor of prosystemin-mediated responses2 gene encodes a fatty acid desaturase required for the biosynthesis of jasmonic acid and the production of a systemic wound signal for defense gene expression. *Plant Cell*. 2003;**15**:1646–61
- Li J, Li B, Luo L. et al. Increased phenolic acid and tanshinone production and transcriptional responses of biosynthetic genes in hairy root cultures of *Salvia przewalskii* Maxim. treated with methyl jasmonate and salicylic acid. *Mol Biol Rep*. 2020;**47**:8565–78
- Li JF, Zhou HP, Zhang Y. et al. The GSK3-like kinase BIN2s a molecular switch between the salt stress response and growth recovery in *Arabidopsis thaliana*. *Dev Cell*. 2020;**55**:367–380.e6
- Li JM, Hu LP, Zhang L. et al. Exogenous spermidine is enhancing tomato tolerance to salinity-alkalinity stress by regulating chloroplast antioxidant system and chlorophyll metabolism. *BMC Plant Biol*. 2015;**15**:303

34. Liu C, Atanasov KE, Arafaty N. et al. Putrescine elicits ROS-dependent activation of the salicylic acid pathway in *Arabidopsis thaliana*. *Plant Cell Environ.* 2020;**43**:2755–68
35. Liu MH, Wei JW, Liu W. et al. S-nitrosylation of ACO homolog 4 improves ethylene synthesis and salt tolerance in tomato. *New Phytol.* 2023;**239**:159–73
36. Liu SH, Zhang PY, Li CC. et al. The moss jasmonate ZIM-domain protein PnJAZ1 confers salinity tolerance via crosstalk with the abscisic acid signaling pathway. *Plant Sci.* 2023;**280**:1–11
37. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods.* 2001;**25**:402–8
38. Lu KK, Song RF, Guo JX. et al. CycC1;1-WRKY75 complex-mediated transcriptional regulation of SOS1 controls salt stress tolerance in *Arabidopsis*. *Plant Cell.* 2023;**35**:2570–91
39. Luo XF, Dai YJ, Zheng C. et al. The ABI4-RbohD/VTC2 regulatory module promotes reactive oxygen species (ROS) accumulation to decrease seed germination under salinity stress. *New Phytol.* 2021;**229**:950–62
40. Ma Y, Xue H, Zhang F. et al. The miR156/SPL module regulates apple salt stress tolerance by activating MdWRKY100 expression. *Plant Biotechnol J.* 2021;**19**:311–23
41. Masmoudi F, Tounsi S, Dunlap CA. et al. Endophytic halotolerant *Bacillus velezensis* FMH2 alleviates salt stress on tomato plants by improving plant growth and altering physiological and antioxidant responses. *Plant Physiol Biochem.* 2021;**165**: 217–27
42. Min DD, Zhou JX, Li JZ. et al. SIMYC2 targeted regulation of polyamines biosynthesis contributes to methyl jasmonate-induced chilling tolerance in tomato fruit. *Postharvest Biol Technol.* 2021;**174**:111443
43. Negi N, Khurana P. A salicylic acid inducible mulberry WRKY transcription factor, MiWRKY53 is involved in plant defence response. *Plant Cell Rep.* 2021;**40**:2151–71
44. Nie W, Liu L, Chen Y. et al. Identification of the regulatory role of SlWRKYs in tomato defense against *Meloidogyne incognita*. *Plants (Basel).* 2023;**12**:2416
45. Ollás R, Eljakaoui Z, Li J. et al. The plasma membrane Na⁺/H⁺ antiporter SOS1 is essential for salt tolerance in tomato and affects the partitioning of Na⁺ between plant organs. *Plant Cell Environment.* 2009;**32**:904–16
46. Prerostova S, Dobrev PI, Gaudinova A. et al. Hormonal dynamics during salt stress responses of salt-sensitive *Arabidopsis thaliana* and salt tolerant *Thellungiella salsuginea*. *Plant Sci.* 2017;**264**: 188–98
47. Qiu ZB, Guo JL, Zhu AJ. et al. Exogenous jasmonic acid can enhance tolerance of wheat seedlings to salt stress. *Ecotoxicol Environ Saf.* 2014;**104**:202–8
48. Raffener M, Üstün S, Guerra T. et al. The *Xanthomonas* type-III effector XopS stabilizes CaWRKY40a to regulate defense responses and stomatal immunity in pepper (*Capsicum annuum*). *Plant Cell.* 2022;**34**:1684–708
49. Rushton PJ, Somssich IE, Ringler P. et al. WRKY transcription factors. *Trends Plant Sci.* 2010;**15**:247–58
50. Schaller F. Enzymes of the biosynthesis of octadecanoid-derived signaling molecules. *J Exp Bot.* 2001;**52**:11–23
51. Shahzad A, Pitann B, Ali H. et al. Maize genotypes differing in salt resistance vary in jasmonic acid accumulation during the first phase of salt stress. *J Agron Crop Sci.* 2015;**201**:443–51
52. Singh D, Debnath P, Sane AP. et al. Tomato (*Solanum lycopersicum*) WRKY23 enhances salt and osmotic stress tolerance by modulating the ethylene and auxin pathways in transgenic *Arabidopsis*. *Plant Physiol Biochem.* 2023;**195**:330–40
53. Sun XC, Gao YF, Li HR. et al. Over-expression of SlWRKY39 leads to enhanced resistance to multiple stress factors in tomato. *J Plant Biol.* 2015;**58**:52–60
54. Tavallali V, Karimi S. Methyl jasmonate enhances salt tolerance of almond rootstocks by regulating endogenous phytohormones, antioxidant activity and gas-exchange. *Plant Physiol.* 2019;**234**:98–105
55. Vanderauwera S, Vandenbroucke K, Inzé A. et al. AtWRKY15 perturbation abolishes the mitochondrial stress response that steers osmotic stress tolerance in *Arabidopsis*. *Proc Natl Acad Sci USA.* 2012;**109**:20113–8
56. Wang PJ, Xu ZJ, Zhang Y. et al. Over-expression of spermidine synthase 2 (SlSPDS2) in tomato plants improves saline-alkali stress tolerance by increasing endogenous polyamines content to regulate antioxidant enzyme system and ionic homeostasis. *Plant Physiol Biochem.* 2022;**192**:172–85
57. Wang X, Ajab Z, Liu CX. et al. Overexpression of transcription factor SlWRKY28 improved the tolerance of *Populus davidiana* × *P. bolleana* to alkaline salt stress. *BMC Genet.* 2020;**21**:103
58. Wang Z, Hong YC, Zhu GT. et al. Loss of salt tolerance during tomato domestication conferred by variation in a Na⁺/K⁺ transporter. *EMBO J.* 2020;**39**:e103256
59. Wasternack C, Song SS. Jasmonates: biosynthesis, metabolism, and signaling by proteins activating and repressing transcription. *J Exp Bot.* 2017;**68**:1303–21
60. Wu J, Wang L, Baldwin IT. Methyl jasmonate-elicited herbivore resistance: does MeJA function as a signal without being hydrolyzed to JA? *Planta.* 2008;**227**:1161–8
61. Wu TY, Goh HZ, Azodi CB. et al. Evolutionarily conserved hierarchical gene regulatory networks for plant salt stress response. *Nature Plants.* 2021;**7**:787–99
62. Xie T, Chen CJ, Li C. et al. Genome-wide investigation of WRKY gene family in pineapple: evolution and expression profiles during development and stress. *BMC genomics.* 2018; **19**: 490.
63. Xu JW, Yang JY, Xu ZJ. et al. Exogenous spermine-induced expression of SlSPMS gene improves salinity-alkalinity stress tolerance by regulating the antioxidant enzyme system and ion homeostasis in tomato. *Plant Physiol Biochem.* 2020;**157**:79–92
64. Xu JJ, Liu T, Yang SC. et al. Polyamines are involved in GABA-regulated salinity-alkalinity stress tolerance in muskmelon. *Environ Exp Bot.* 2019;**164**:181–9
65. Xu ZJ, Wang F, Ma YB. et al. Transcription factor SlAREB1 is involved in the antioxidant regulation under saline-alkaline stress in tomato. *Antioxidants.* 2022;**11**:1673
66. Xu ZJ, Wang JC, Ma YB. et al. The bZIP transcription factor SlAREB1 regulates anthocyanin biosynthesis in response to low temperature in tomato. *Plant J.* 2023;**115**:205–19
67. Yang JY, Wang PJ, Li SZ. et al. Polyamine oxidase triggers H₂O₂-mediated spermidine improved oxidative stress tolerance of tomato seedlings subjected to saline-alkaline stress. *Int J Mol Sci.* 2019;**23**:1625
68. Yang JY, Wang PJ, Li XJ. et al. Regulation of cold resistance by the polyamine biosynthetic gene SlSPDS2 via modulating the antioxidant enzyme system and osmotic regulatory substances in *Solanum lycopersicum*. *Environ Exp Bot.* 2023;**216**: 105531
69. Yu ZP, Duan XB, Luo L. et al. How plant hormones mediate salt stress responses. *Trends Plant Sci.* 2020;**25**:1117–30
70. Yu J, Zhu CS, Xuan W. et al. Genome-wide association studies identify OsWRKY53 as a key regulator of salt tolerance in rice. *Nat Commun.* 2023;**14**:3550
71. Zhang H, Zhang Q, Zhai H. et al. Transcript profile analysis reveals important roles of jasmonic acid signaling pathway in

- the response of sweet potato to salt stress. *Sci Rep.* 2017;**7**:40819
72. Zhang LL, Zheng Y, Xiong XX. et al. The wheat VQ motif-containing protein *TaVQ4-D* positively regulates drought tolerance in transgenic plants. *J Exp Bot.* 2023;**74**:5591–605
73. Zhang X, Chen LC, Shi QH. et al. *SlMYB102*, an R2R3-type MYB gene, confers salt tolerance in transgenic tomato. *Plant Sci.* 2020;**291**:110356
74. Zhao CZ, Jiang W, Zayed O. et al. The LRXs-RALFs-FER module controls plant growth and salt stress responses by modulating multiple plant hormones. *Natl Sci Rev.* 2021;**8**:nwaa149.
75. Zhao Y, Dong W, Zhang NB. et al. A wheat allene oxide cyclase gene enhances salinity tolerance via jasmonate signaling. *Plant Physiol.* 2014;**164**:1068–76
76. Zhou J, Wang J, Zheng ZY. et al. Characterization of the promoter and extended C-terminal domain of *Arabidopsis WRKY33* and functional analysis of tomato *WRKY33* homologues in plant stress responses. *J Exp Bot.* 2015;**66**:4567–83